

REVIEW

Phosphoinositide-derived messengers in endocrine signaling

T Balla

Endocrinology and Reproduction Research Branch, NICHD, National Institutes of Health, Building 49, Room 6A35, 49 Convent Drive, Bethesda, Maryland 20892, USA

(Requests for offprints should be addressed to T Balla; Email: ballat@mail.nih.gov)

Abstract

One of the fundamental questions in endocrinology is how circulating or locally produced hormones affect target cell functions by activating specific receptors linked to numerous signal-transduction pathways. An important subset of G protein-coupled cell-surface receptors can activate phospholipase C enzymes to hydrolyze a small but critically important class of phospholipids, the phosphoinositides. Although this signaling pathway has been extensively explored over the last 20 years, this has proven to be only the tip of the iceberg, and the multiplicity and diversity of the cellular functions controlled by phosphoinositides have surpassed any imagination. Phosphoinositides have been found to be key regulators of ion

channels and transporters, and controllers of vesicular trafficking and the transport of lipids between intracellular membranes. Essentially, they organize the recruitment and regulation of signaling protein complexes in specific membrane compartments. While many of these processes have been classically studied by cell biologists, molecular endocrinology cannot ignore these recent advances, and now needs to integrate the cell biologist's views in the modern concept of how hormones affect cell functions and how derailment of simple molecular events can lead to complex endocrine and metabolic disorders.

Journal of Endocrinology (2006) **188**, 135–153

Introduction

The mechanism of action of 'Ca²⁺-mobilizing hormones', a term introduced in the mid-1970s to denote hormones that exert their effects through generation of a cytosolic Ca²⁺ signal, was largely elucidated by the end of the 1980s (Fig. 1 and 2). The basic elements of this canonical signal-transduction cascade include the activation of the phospholipase C (PLC) β and γ isoforms by G protein-coupled receptors (GPCRs) and receptor tyrosine kinases (RTKs) respectively. Stimulation of these receptors generates two second messengers, the soluble Ins(1,4,5)P₃ (InsP₃) and the hydrophobic diacylglycerol from membrane, PtdIns(4,5)P₂ (Berridge 1984). Once generated, the Ca²⁺-mobilizing messenger, InsP₃, binds to specific receptors that are located mainly in the endoplasmic reticulum (ER) and function as tetrameric cation channels to release Ca²⁺ from intracellular Ca²⁺ stores (Mikoshiba 1997). This endogenous Ca²⁺ signal, in combination with increased Ca²⁺ influx via multiple Ca²⁺ entry mechanisms, activates numerous cytoplasmic and membrane-bound effector molecules with the help of Ca²⁺-binding proteins. Ca²⁺ is also taken up by mitochondria, where it

stimulates metabolic enzymes (Hajnoczky *et al.* 1995) or initiates complex responses such as apoptosis (Pacher & Hajnoczky 2001). The other limb of this messenger system, diacylglycerol (DAG), directly activates several members of the protein kinase C family (Nishizuka 1988), and can also contribute to the direct regulation of some ion channels (Hardie 2003).

This elegantly simple concept marked only the beginning of an amazing explosion of research into phosphoinositides, and almost every component of the calcium-phosphoinositide messenger system has grown to become a research field of its own. The metabolism of InsP₃ via both phosphorylation and sequential dephosphorylation revealed the identity of several inositol phosphate isomers (Shears 1998), as well as the InsP kinases (Verbsky *et al.* 2005) and phosphatases (Majerus *et al.* 1999). We are just at the beginning of understanding which of the numerous inositol phosphate isomers have regulatory functions. The discovery of phosphoinositide 3-kinases (PI3K) was the starting point in the recognition of the function of phosphoinositides as membrane-bound signaling molecules and was one of the main driving forces behind the search for proteins that are downstream targets




Figure 1 Updated version of the classical 'PI cycle'. Plasma membrane PtdIns(4,5)P₂ is hydrolyzed by PLC enzymes, after receptor stimulation, to generate two second messengers, Ins(1,4,5)P₃ and DAG. Ins(1,4,5)P₃, which mobilizes intracellular Ca²⁺, is rapidly degraded by sequential dephosphorylations to yield myo-inositol. (A fraction of the Ins(1,4,5)P₃ is converted to higher inositol phosphates via pathways that differ between plants, yeast and vertebrates (Shears 2004), but it is not shown in the figure for better clarity.) Some of the dephosphorylating enzymes on the pathway of Ins(1,4,5)P₃ degradation are sensitive to inhibition by Li⁺ ions, a finding that led to the idea that the therapeutic effects of Li⁺ in bipolar disorder might be related to altered signaling affecting primarily hyperactive receptors (Berridge *et al.* 1989). DAG, the other product of PtdIns(4,5)P₂ hydrolysis, activates PKC enzymes (and some other effectors, such as ion channels or protein kinase D (PKD) – not shown) before being converted to PtdA by DAG-kinase enzymes. PtdA of the plasma membrane can also activate certain effectors (not shown). In the endoplasmic reticulum (ER), PtdA is converted to CDP-DAG, which is conjugated with myo-inositol by PI synthase enzyme(s). When cells are stimulated in the presence of Li⁺, CDP-DAG accumulates because of the shortage of myo-inositol derived from phosphoinositide hydrolysis. The PtdIns of the ER is transferred to the plasma membrane by PI transfer proteins, but it is less clear whether a similar mechanism transfers PtdA between the membranes. Most downstream pathways in this messenger system are regulated by PKC and other Ca²⁺-responsive protein kinases or phosphatases.

of the inositol phospholipids. With the identification of new forms of inositol lipids and newly discovered functions of their classical forms, came the cloning of the multiple forms of inositol lipid kinases (Fruman *et al.* 1998) and phosphatases (Majerus *et al.* 1999), several of which have also been linked to human diseases (Pendaries *et al.* 2003) (Fig. 3). Research into the calcium aspect of InsP₃-mediated signaling has also shown an enormous expansion, starting with the cloning of the InsP₃ receptor (Furuichi *et al.* 1989, Mignery *et al.* 1989) and the continuing quest to identify the molecular entities behind the store-operated Ca²⁺ entry (SOC) pathway (Parekh & Putney 2005). There are several excellent reviews covering almost every aspect of inositol research from the viewpoint of the protein classes as indicated above. This review cannot be a compilation of all of these areas. Instead, it will aim to

summarize recent advances and open questions from selected areas of endocrine research pointing to the inositol lipid angle within those areas. This is a selection with an inevitable bias and is by no means an indication that other aspects of inositides are not important in endocrinology. After all, it could be claimed that there are only two kinds of research: one in which phosphoinositides are already implicated and another where their involvement is yet to be recognized!

Receptor-mediated InsP₃ formation and Ca²⁺ release

When Ca²⁺-mobilizing receptors engage their agonist ligands, the subsequent activation of PLC enzymes at the plasma membrane leads to rapid breakdown of




Figure 2 G protein-coupled receptors (GPCRs) and receptor tyrosine kinases (RTKs) activate PtdIns(4,5)P₂ hydrolysis by distinct mechanisms, the former using G protein-mediated activation of PLCβ isoforms, and the latter causing PLCγ activation. Ins(1,4,5)P₃ binds to its receptors located primarily in the ER in the form of tetrameric Ca²⁺ channels releasing Ca²⁺ from the internal stores. The N-terminal Ins(1,4,5)P₃-binding domain of the InsP₃ receptor is probably in close proximity to the C-terminal channel domain with a long intervening regulatory domain (see insert). Emptying of the ER Ca²⁺ pools relays information (by unknown mechanisms) to plasma membrane Ca²⁺ channels, commonly termed store-operated Ca²⁺ channels (SOC), to enhance Ca²⁺ influx into the cell. Members of the TRP family of Ca²⁺ channels may function as SOC, and it has been suggested that there is a physical regulatory contact between plasma membrane-adjacent InsP₃ receptors and certain TRP channels (see text for details). Another spatial organization is related to the ‘coupling’ of Ca²⁺ release via the InsP₃ receptors with uptake by mitochondria that are juxtaposed to InsP₃ receptor-rich ER domains.

PtdIns(4,5)P₂. This process is both amplified and at the same time self-limited in a number of ways. It is amplified because of the positive regulation of PLC enzymes by cytoplasmic Ca²⁺, an effect that accounts for the profound impact of Ca²⁺ release on the extent of PtdIns(4,5)P₂ hydrolysis (e.g. Horowitz *et al.* 2005). Without receptor activation, Ca²⁺ per se is much less potent, and only large cytoplasmic Ca²⁺ increases lead to PtdIns(4,5)P₂ breakdown. Although all PLC enzymes are Ca²⁺-sensitive, the PLCδ1 enzyme appears to be largely responsible for Ca²⁺-mediated positive feedback on InsP₃ generation (Rhee 2001). At the same time, the process of InsP₃ formation is limited because most GPCRs undergo

homologous desensitization that limits their ability to activate G-proteins (Luttrell & Lefkowitz 2002). Moreover, both GPCRs and RTKs undergo ligand-induced endocytosis and are rapidly removed from the plasma membrane, whence they can also be quickly recycled and reappear during resensitization (Gaborik & Hunyady 2004). PtdIns(4,5)P₂ hydrolysis is also limited because, as a result of its relatively low level in the membrane, it has to be replenished by the sequential actions of PI 4- and PIP 5-kinases on the larger pools of PtdIns. Even the larger pool of plasma membrane PtdIns can become depleted if not maintained by supply from the ER by PI transfer proteins (Fig. 1).




Figure 3 The network of interconversions between phosphoinositides. Phosphoinositides are phosphorylated (red arrows) by inositol kinases and are dephosphorylated (blue arrows) by phosphoinositide phosphatases. Most (but not all) of the kinases are quite specific and phosphorylate only one substrate in a specific position on the inositol ring. Some kinases (such as the class II PI 3-kinases, PI3KC2) show wider substrate tolerance *in vitro* but are likely to have a clear substrate preference *in vivo*. The substrate specificity of most (but not all) phosphatases is less stringent *in vitro*, and some can dephosphorylate lipids as well as the soluble inositol phosphates. The physiologic and preferred substrate is not known for all inositol phosphatases, nor is it known in all cases which enzyme or enzymes are primarily responsible for a specific interconversion step. For example, Sac1 phosphatases can remove the 3- and 4-phosphates from a mono-phosphorylated inositol lipid, but synaptojanins (Snj), which are primarily known as 5-phosphatases, also contain a Sac1 phosphatase motif and can function as Sac1 phosphatases (Guo *et al.* 1999). Since myotubularin-related inositol lipid phosphatases (MTMRs) (Taylor & Dixon 2003) also dephosphorylate the 3 position on PtdIns(3)P, it is likely that PtdIns(3)P is dephosphorylated by Sac1, Snj-s and MTMRs in different cellular compartments (Parrish *et al.* 2004). This figure indicates only the more established routes; for all other possibilities of interconversion, see Abel *et al.* (2001).

Which inositol lipid pools participate in the classical signaling cascade and which enzymes are primarily responsible for their generation?

The existence of hormone-sensitive and insensitive pools of phosphoinositides has been described in early research using metabolic labeling with myo-inositol (Koreh & Monaco 1986). Recent studies have indicated that PtdIns(4,5)P₂ is produced by two different pathways: in the canonical pathway, it is formed through PtdIns(4)P by sequential phosphorylations by PI 4-kinases (Balla 1998)

and the type I PIP kinases (or PIP5Ks) (Hinchliffe *et al.* 1998). In a recently recognized pathway, PtdIns(4,5)P₂ synthesis proceeds via PtdIns(5)P that is phosphorylated by type-II PIP kinases (or PIP4Ks) (Rameh *et al.* 1997) (Fig. 3). Since each group of enzymes has multiple forms that are located in distinct cellular compartments, it is inevitable that phosphoinositides will be generated in different types of membranes. Which of these inositol pools are linked to hormone action? It is widely assumed that hormonal stimulation leads to breakdown of the plasma membrane PtdIns(4,5)P₂ pool, and this has been

substantiated by the use of green fluorescent protein (GFP)-fused pleckstrin homology (PH) domains that recognize the lipid in the plasma membrane. However, PtdIns(4,5)P₂ pools also exist in other membrane locations, such as the Golgi, the ER and the nucleus (Watt *et al.* 2002), and PLC enzymes are also found at these sites (Rebecchi & Pentyala 2000). Therefore, it is not unreasonable to assume that PLC activation – either direct, or mediated by the cytoplasmic Ca²⁺ increase – occurs at intracellular membranes and contributes to InsP₃ and DAG generation at those sites with local consequences.

Equally important is the simple question of which enzymes generate the PI(4,5)P₂ that is subject to receptor-controlled PLC-mediated hydrolysis. Recently, it was shown that human PIP 5-kinase γ is the enzyme that is necessary for GPCR-mediated InsP₃ formation and Ca²⁺ signaling (Wang *et al.* 2004). In contrast, mouse PIP 5-kinase I β (identical to human PIP5KI α), recruited by Bruton's tyrosine kinase, enhances Ca²⁺ signaling in B cells (Saito *et al.* 2003). Stimulation of PIP 5-kinase activity by small guanosine triphosphate (GTP)-binding proteins has been well documented (Chong *et al.* 1994, Honda *et al.* 1999), but it is not known whether receptor-mediated activation of PIP 5-kinase activity occurs. A very rapid increase in PtdIns(4,5)P₂ level after stimulation has been recently demonstrated, suggesting direct activation of PtdIns(4,5)P₂ synthesis (Xu *et al.* 2003). Less clear is the identity of the PI 4-kinase or kinases that provide PtdIns(4)P for the PIPKs. Hormone-sensitive PtdIns(4)P pools have been shown to be generated by the type-III PI4Ks (which are sensitive to higher concentrations of PI3K inhibitors) (Nakanishi *et al.* 1995). However, neither the PI4 KIII α nor - β isoform is detectable at the plasma membrane of mammalian cells; instead, these isoforms are located in ER and Golgi membranes respectively (Wong *et al.* 1997). Yet, recent studies have shown that a plasma membrane (PM) pool of PtdIns(4)P is generated by PI4 KIII α (Balla *et al.* 2005), raising the question of whether the lipid is generated at the PM by a small fraction of the enzyme or is generated elsewhere and transported to the PM. There is little direct evidence that PI4K activities are stimulated during receptor activation, and only PI4 KII β has been shown to be activated by a Rac-dependent mechanism (Wei *et al.* 2002).

As mentioned above, the maintenance of PtdIns(4,5)P₂ pools also requires the function(s) of the family of PI transfer proteins (PITPs). These proteins transfer PtdIns and PtdCho between membranes and are necessary for maintenance of the PLC-sensitive phosphoinositide pools (Thomas *et al.* 1993). PITPs also exist in several forms, of which the soluble PITP α and PITP β isoforms have non-overlapping functions that reach far beyond the regulation of hormone-sensitive phosphoinositide pools, affecting both phospholipid synthesis and vesicular trafficking. This function of lipid-transfer proteins will be further detailed below.

How does InsP₃ regulate Ca²⁺ release?

Since the isolation and cloning of the InsP₃ receptor channels, impressive progress has been made in understanding their functions. All three isoforms of the InsP₃ receptor (types I, II and III) function as intracellular Ca²⁺ channels that operate as a homo- or heterotetramer (Mikoshiba 1993). Each receptor subunit has a channel portion containing six transmembrane helices and a pore domain located between TM5 and TM6, close to the C-terminus of the protein (Maeda *et al.* 1991, Galvan *et al.* 1999, Patel *et al.* 1999). The ligand-binding domain (LBD) of the receptor is located at the N-terminus (Mignery & Sudhof 1990) and is separated from the channel domain by a long intervening regulatory region facing the cytoplasm (Mignery & Sudhof 1990, Mikoshiba 1993). InsP₃ binding leads to rapid activation of the channel, but Ca²⁺-induced Ca²⁺ release, similar to that featured in the related ryanodine receptors (RyR), is also an important regulatory mechanism of IP₃Rs (Taylor & Laude 2002). Because of their ER location, little is known about the gating mechanism and properties of IP₃R channels. The only electrophysiologic data are derived from isolated channels reconstituted in lipid bilayers (Bezprozvanny *et al.* 1991) or from patch-clamp recordings of the nuclear envelope (Mak & Foskett 1994). Most of our knowledge of the behavior of the intact channels is inferred from measurements of Ca²⁺ (or other cation) fluxes. These studies have provided invaluable information about the InsP₃ and Ca²⁺ regulation of the channel (Bezprozvanny *et al.* 1991, Hajnoczky & Thomas 1994), but no molecular mechanisms or states have been correlated with channel behavior. One major question is how the N-terminal LBD can regulate the channel itself. Recent evidence suggests that the C-terminal channel domain and the N-terminal LBD are in very close molecular proximity, and that the ligand-induced conformational change within the LBD could be transferred to the channel domain itself (Boehning & Joseph 2000). The recently solved X-ray structure of the LBD (Bosanac *et al.* 2002) and the adjacent N-terminal inhibitory domain (Bosanac *et al.* 2005) has helped to clarify the structural basis of InsP₃ binding, and, together with high-resolution electron microscopy and 3-D reconstruction of the channel structure (Jiang *et al.* 2002, da Fonseca *et al.* 2003, Sato *et al.* 2004), should advance our understanding of the gating mechanism of the protein.

The positioning of ER membranes containing InsP₃ receptors relative to other membranes, and the interaction of the receptor with other proteins, add to the complexity of Ca²⁺ regulation in local compartments (Fig. 2). Early fractionation studies suggested that InsP₃ receptors could be isolated from 'mitochondrial' (Dawson & Irvine 1984) and 'plasma membrane' (Guillemette *et al.* 1988) fractions, indicating ER contamination of these fractions enriched in InsP₃ receptors. Recent evidence has shown the

existence of a special ER-mitochondria interface (Rizzuto *et al.* 1993) and a very close 'quasi-synaptic' functional coupling between InsP_3 receptor-mediated Ca^{2+} release and mitochondrial Ca^{2+} uptake (Csordas *et al.* 1999). Moreover, the physical association of InsP_3 receptors with TRPC3 channels at the plasma membrane could provide the basis for Ca^{2+} influx regulation linked to InsP_3 -induced Ca^{2+} release (Kiselyov *et al.* 1998). These observations raise the question of whether the InsP_3R could serve as a structural participant in the tethering of the ER to other membranes in the proximity of the channel. It has also been suggested that the LBD of the InsP_3R binds to $\text{PtdIns}(4,5)\text{P}_2$ of the plasma membrane in quiescent cells when InsP_3 levels are low (Glouchankova *et al.* 2000), and that changing $\text{PtdIns}(4,5)\text{P}_2$ together with increased InsP_3 could also participate in the regulation of these channels.

Cell-surface receptors and PI3K-mediated signaling

The classical activation mechanism of PI3Ks was identified by studies on receptor tyrosine kinases and on soluble and oncogenic tyrosine kinases (Otsu *et al.* 1991). In these systems, the p85 regulatory subunit associates with tyrosine-phosphorylated target sequences through its SH2 domains, recruiting the cytoplasmic PI3K α or β catalytic subunit to the membrane. The p85 subunit becomes tyrosine phosphorylated during this process, leading to increased activity of the kinase. Less is known about the mechanism of PI3K activation in the case of GPCRs. In hematopoietic cells, where PI3K γ is found in significant amounts, activation via the $\beta\gamma$ -subunits of Gi/Go proteins is the main activation pathway (Stephens *et al.* 1994). This is mediated by associated p101 (Stephens *et al.* 1997) or p84 (Suire *et al.* 2005) regulatory subunits, but direct regulation of the PI3K γ enzyme by $\beta\gamma$ -subunits has also been reported (Leopoldt *et al.* 1998). Much less clear and more controversial is the manner in which PI3K is activated by GPCRs in tissues where PI3K γ is not expressed, or is present only at low levels. In many cases, activation occurs by transactivation of receptor tyrosine kinases (Daub *et al.* 1996) followed by the above-detailed mechanism, but this is not the sole means by which GPCRs activate PI3Ks. It is also not known what determines which of the class I PI3Ks (α, β, δ) is activated and which splice form of the p85/p55 regulatory subunit associates with them. Deletion of either PI3K α or β is lethal (Bi *et al.* 1999, 2002), as is the elimination of all the splice forms of the p85/55 α subunits (Fruman *et al.* 2000), but not of the p85 α form alone (Terauchi *et al.* 1999), indicating a level of complexity that is still far from being understood. Impairment of PI3K signaling has prominent effects on insulin signaling (Terauchi *et al.* 1999), and recent studies indicate that the stoichiometry between

the p85 and p110 subunits, together with the direct interaction of the free p85 subunit with IRS-1, is a major factor determining the insulin responsiveness of the cells (Luo *et al.* 2005).

Activation of PI3K by estrogens via ligand-dependent association of ER α with the p85 regulatory subunit has been demonstrated (Simoncini *et al.* 2000). More recently, a heptahelical receptor, GPR30, which is located in the ER, was shown to respond to estrogen stimulation, leading to the activation of PI3K in the nucleus, as indicated by the translocation of the Akt PH-domain to the nucleus (Revankar *et al.* 2005). Many questions remain to be answered about this intriguing effect, but it appears to be a new paradigm in steroid hormone action that is not mediated by the classical nuclear receptors.

The main downstream signaling pathway from PI3Ks proceeds via the Akt protein kinase (Franke *et al.* 1997) and its upstream regulator kinase, PDK1 (Alessi *et al.* 1997). These kinases are master regulators of a whole range of cellular processes related to glucose metabolism, protein synthesis and cell division, and also represent the main antiapoptotic pathway. Detailed coverage of these processes in endocrine and metabolic regulation is beyond the scope of this review, but can be found elsewhere (Mora *et al.* 2004).

Receptor trafficking and its regulation by phosphoinositides

Many cell-surface receptors undergo ligand-induced endocytosis mostly (but not exclusively) by a clathrin-mediated internalization process that shares many of the characteristics of the endocytosis and recycling of nutrient receptors (Brown & Goldstein 1979). The sorting of the receptors into clathrin-coated pits is mediated by interaction of the receptor with clathrin-adaptor proteins. Several adapter proteins have been identified, including the tetrameric adapters, AP-(2-4), the monomeric adapters such as AP-180 (or its nonneural form, CALM), the Dab1/ARH and the GGA proteins. Many of these adapters exert their effects at intracellular membranes (Owen *et al.* 2004), when they recognize specific sequences (sorting motifs) within the intracellular segments of nutrient receptors or RTKs. They also bind to clathrin, thereby bringing the receptors to the site of clathrin assembly (Fig. 4). Several sorting motifs have been identified in receptors, some containing Tyr, such as the NPxY or the Yxx Φ sequences, and others containing Leu/Ile residues, such as the D/ExxxL/I or DxxLL motifs (Bonifacino & Traub 2003). In the case of GPCRs, the common adapter proteins are β -arrestins which bind to GPCR tails that are Ser/Thr phosphorylated by G protein receptor kinases (GRKs) (Lefkowitz 1993). Phosphorylation of GPCR tails by GRKs is greatly facilitated by the binding of agonist ligands, and it accounts for the ligand-dependence of GPCR internalization. The role of receptor internalization is obvious in the case of nutrient




Figure 4 The general scheme of endocytosis from the plasma membrane. Transmembrane proteins (e.g. receptors) associate with adaptor proteins capable of recognizing the internalization signal (usually within their membrane-adjacent cytoplasmic sequence). (Often, a covalent modification, such as phosphorylation or ubiquitination, takes place at these sites on the transmembrane proteins – not shown here.) The adaptor protein interaction is also aided by the presence of membrane phosphoinositides and requires an active small GTP-binding protein (such as Arfs or Rabs). The adaptor protein also binds clathrin, thereby recruiting the protein into clathrin-coated pits. Several adaptor proteins have been shown to bind phosphoinositides, which, in most cases, are the most abundant PtdIns(4,5)P₂. Once the vesicles are pinched off (clathrin-coated vesicles, CV), this lipid is no longer demonstrable in the vesicles and is replaced by 3-phosphorylated inositides, contributing to a new identity of the vesicles. Recent studies also indicate that PtdIns(3,4,5)P₃ formed in the membrane can be a source of 3-phosphorylated lipids in the endosomes after sequential dephosphorylations by vesicle-associated phosphatases that remove the 5- and 4- phosphates (Ivetac *et al.* 2005, Shin *et al.* 2005). Multivesicular bodies are formed from early endosomes (EE) after inward invagination and budding, and PtdIns(3,5)P₂ is a critical component of this process (see text for more details).

receptors but is less clear in the case of hormone receptors. For this latter, it is an important mechanism for regulating the number of available receptors on the cell surface, and it also eliminates activated receptors that often undergo degradation (Dikic 2003). There is also increasing evidence that some internalized receptors may continue to signal in the endocytic compartments (Luttrell & Lefkowitz 2002).

It is quite remarkable that many of the proteins that participate in the internalization process contain binding sites for phosphoinositides. With the exception of the GGAs, all of the above-mentioned clathrin-adaptor proteins have been shown to contain one or more

phosphoinositide-binding motifs (Owen *et al.* 2004). Some GRKs (such as GRK2 and 3) contain a pleckstrin-homology domain that is important for the membrane recruitment of the enzymes (Pitcher *et al.* 1998), but the phospholipid species involved in their regulation is not yet known (Carman *et al.* 2000). Recently, β -arrestins were shown to contain phosphoinositide-binding regions that are important for GPCR internalization (Gaidarov *et al.* 1999, Lee *et al.* 2003). Finally, the GTP-binding protein, dynamin, which is critical for pinching off internalized vesicles from the plasma membrane, is also known to contain a PH domain. In this case, mutation of the residues essential for lipid binding has a dominant negative

effect on receptor endocytosis (e.g. Lee *et al.* 1999). From the presence of these domains in the above proteins and their ability to bind phosphoinositides *in vitro*, it is widely assumed that phosphoinositides regulate receptor endocytosis. Mutagenesis studies targeting the lipid-binding regions of the proteins support this assumption, but few studies are available on the question of whether modifying lipid production would affect receptor trafficking. Wortmannin-sensitive (i.e., type III PI4K-synthesized) PtdIns(4)P and PtdIns(4,5)P₂ pools support muscarinic and β 2-adrenergic receptor endocytosis (Sorensen *et al.* 1998), and the plasma membrane recruitment of AP-2 protein and transferrin receptor endocytosis correlates with PIP5K1 β activity (Padron *et al.* 2003). In general, it is not certain which phosphoinositide species regulates any particular step in the complex process of endocytosis. Because of its abundance and *in vitro* binding to many of the above domains, PtdIns(4,5)P₂ is considered the most important inositide in receptor endocytosis. Although several reports have indicated that PI 3-kinases are also agents in this process (Joly *et al.* 1994, Naga Prasad *et al.* 2002), pharmacologic blockade of PI3Ks does not dramatically influence the early steps of receptor internalization (Hunyady *et al.* 2002). Most recently, the protein kinase activity of PI3K γ via phosphorylation of tropomyosin was shown to be critical for β -adrenergic receptor endocytosis (Naga Prasad *et al.* 2005).

Fate of the internalized receptors

It is also well documented that the subsequent fate of internalized receptors is highly dependent on processes regulated by PI3Ks. After reaching early endosomes, a large proportion of receptors recycle back to the plasma membrane via a vesicular mechanism involving Rab4 and Rab5 GTPases (Seachrist *et al.* 2000, Hunyady *et al.* 2002, Dale *et al.* 2004). This rapid recycling process is inhibited by the PI 3-kinase inhibitors, wortmannin and LY294002, leading to the accumulation of the receptors in large endocytic vesicles (Shpetner *et al.* 1996, Hunyady *et al.* 2002). A fraction of the transferrin receptors and some GPCRs are also sorted into another compartment, called recycling endosomes, that is positive for Rab11 GTPase, and from which there is a significantly slower recycling to the cell surface by a process more resistant to PI 3-kinase inhibitors (Hunyady *et al.* 2002, van Dam *et al.* 2002). Interestingly, a group of GPCRs very rapidly recycle back to the cell surface after internalization, and these receptors are not sorted into, and do not take β -arrestin, to 'deeper' compartments (Zhang *et al.* 1999). One of the most exciting 'organelles' of the internalization and recycling pathway is the multivesicular body (MVB), a site for molecular decisions on whether receptors recycle or undergo degradation. Ubiquitination of receptors often determines their fate. Mono-ubiquitination (sometimes at multiple sites) has been shown to be important for

internalization and targeting of activated RTKs, such as the EGFR to the inner membrane of the MVB and subsequent lysosomal degradation (Hicke 2001, Katzmann *et al.* 2002, Haglund *et al.* 2003). This process is to be distinguished from poly-ubiquitination (where the added ubiquitin is further ubiquitinated multiple times) of soluble proteins that are targeted for proteasomal degradation (Bonifacino & Weissman 1998), although the distinction between the two kinds of ubiquitination is not always clear. Unlike transferrin receptors, some GPCRs also appear at the MVB during their recycling or degradation (Hunyady *et al.* 2002), and also have been shown to be ubiquitinated (Marchese & Benovic 2001, Shenoy *et al.* 2001). The role and significance of GPCR ubiquitination in the endocytic and recycling process is not as well understood as for RTKs. 3-Phosphorylated inositides are principal regulators of the sorting process along the endocytic pathway. Class III PI3Ks generate PtdIns(3)P, which contributes to the recruitment to endocytic vesicles of proteins containing FYVE, or PX domains such as Hrs or sorting nexins, respectively. These proteins are important in cargo selection and vesicle dynamics. PtdIns(3)P is also converted to PtdIns(3,5)P₂ by the PIKfyve enzyme (Shisheva *et al.* 1999) (also termed type III PIP kinase). In *Saccharomyces cerevisiae*, this kinase, termed Fab1p, is also needed for protein sorting into the MVB (Odorozzi *et al.* 1998).

These selected examples illustrate the complexity of the phosphoinositide requirement of the endocytic and sorting process, and highlight the magnitude of the task that remains to be completed in clarifying the molecular details and biologic importance of receptor endocytosis in endocrine functions.

Phosphoinositides and ion channels

The relationship between phosphoinositide turnover and Ca²⁺ signaling has long been firmly established, and for a while the only debate was about which was the consequence of the other (Cockcroft 1981, Michell 1982). However, the discovery of InsP₃-mediated Ca²⁺ release from internal pools has clarified the primary connection between them. Importantly, Ca²⁺ also enters the cells via Ca²⁺ channels located in the plasma membrane, and initially a distinction was made between voltage-gated Ca²⁺ channels, which mostly regulate the functions of 'excitable' cells, and non-voltage-gated Ca²⁺ influx pathways (or receptor-regulated Ca²⁺ channels) working in nonexcitable and endocrine cells. Although this distinction has not persisted very long (e.g., the presence of voltage-gated Ca²⁺ channels in endocrine cells has become common knowledge), the molecular entities behind the Ca²⁺ entry mechanism(s), described as 'capacitative' Ca²⁺ entry pathway or store-operated Ca²⁺ entry (SOC) (Putney 1986), has remained elusive.

All nonexcitable cells display increased Ca^{2+} influx that falls within the criteria of capacitative or SOC, after activation by a 'calcium-mobilizing' stimulus (Putney 1986). After decades of studies to identify the Ca^{2+} channels that underlie this phenomenon, it is still questionable whether a single molecular entity is responsible for the enhanced Ca^{2+} influx observed after emptying of the intracellular Ca^{2+} stores (Parekh & Putney 2005). The current activated by the release of Ca^{2+} stores is termed I_{CRAC} (calcium release-activated current) (Hoth & Penner 1992), but its molecular equivalent is still unresolved, as is the question of whether I_{CRAC} is 'the current' corresponding to SOC (see Parekh & Putney (2005) for an excellent recent review). The Ca^{2+} channel underlying the *Drosophila* transient receptor potential (Trp) in the fly photoreceptor was proposed to be a channel corresponding to SOC (Wes *et al.* 1995). After cloning of the mammalian homologs of these proteins (TrpC₁₋₇, 'C' stands for classical), a large number of studies addressed this question. Of the seven TrpC channels, most data were gathered for TrpC1 and TrpC3 channels. Overexpression of TrpC1 channels yields a nonselective Ca^{2+} -permeable conductance that moderately increases Ca^{2+} influx after store depletion in some studies (Zhu *et al.* 1996), but not in others (Lintschinger *et al.* 2000). This discrepancy could be due to the fact that TrpC1 proteins require coexpression of other Trp channels (such as TrpC4 and -5) to traffic properly to the plasma membrane (Strubing *et al.* 2001). Downregulation or elimination of TrpC1 channels at the cellular level also indicates that they are at least partially responsible for store-operated Ca^{2+} influx (Liu *et al.* 2003) and I_{CRAC} (Mori *et al.* 2002).

Studies on TrpC3 channels provided additional insights into the connection between Ca^{2+} release and influx. Early studies suggested that these channels are regulated by Ca^{2+} and diacylglycerol (and thereby responded to agonist stimulation), but not by store depletion (Lintschinger *et al.* 2000). However, other studies have shown that TrpC3 channels do respond to store depletion and that they physically interact with InsP_3 receptors, providing experimental support for the previously hypothetical conformational-coupling model of SOC (Kiselyov *et al.* 1998). It is very likely that Trp channel behavior depends on the presence of other Trp channels (or of associated proteins) in a particular cell, and that cells display a great variety of Ca^{2+} influx characteristics, depending on the composition of the molecular complex responsible for the store-operated influx phenomenon in specific cell types. This may explain why overexpression studies with the various Trp channels yield apparently discordant information. A detailed, up-to-date review of all Trp channels is given by Putney (2004). Very recent data identify STIM1, an ER-resident, single-transmembrane protein with a Ca^{2+} -sensing luminal domain, as a critical component of SOC (Liou *et al.* 2005, Roos *et al.* 2005). STIM1 is not a channel itself, but it may finally provide us with

a thread from which the SOC phenomenon will be deciphered. There is very little information about direct regulation of TrpC channels by phosphoinositides, but findings with members of other Trp channel families (see below) indicate that this question should be further investigated.

Curiously, the first ion channels for which phosphoinositide regulation was described were not Ca^{2+} channels, but members of the inwardly rectifying (Kir) potassium channel family (Hilgemann & Ball 1996). Some Kir channels were shown to be regulated by the $\beta\gamma$ -subunits of heterotrimeric G proteins (GIRKs) (Krapivinsky *et al.* 1995) and $\text{PtdIns}(4,5)\text{P}_2$ (Huang *et al.* 1998, Zhang *et al.* 1999) and became the prototypical examples. A number of studies followed that showed phosphoinositide regulation of other ion channels. These included other potassium channels, such as the M-current (Suh & Hille 2002) and the underlying KCNQ channels (Zhang *et al.* 2003), and the two-pore domain K^+ channels (Chemin *et al.* 2005, Lopes *et al.* 2005). Among Ca^{2+} channels, members of the Trp family (other than TrpCs) have been shown to be regulated by $\text{PtdIns}(4,5)\text{P}_2$. As shown for the TrpM5, -7 and -8 channels, $\text{PtdIns}(4,5)\text{P}_2$ is necessary for channel activity (Runnels *et al.* 2002). In the case of TrpM8, both cooling and addition of activator ligands, such as menthol, alter the lipid affinity of the channel. The region responsible for lipid regulation has been mapped within the so-called Trp-domain (Rohacs *et al.* 2005). Other Trp channels, such as the vanilloid receptor (TrpV1), are also regulated by $\text{PtdIns}(4,5)\text{P}_2$ (Chuang *et al.* 2001), and this lipid was found to be essential for the recovery of the TrpV1 channel from desensitization (Liu *et al.* 2005). The intimate relationship between some of these channel proteins and PLC enzymes strongly suggests that the Ca^{2+} that enters via Trp channels controls local $\text{PtdIns}(4,5)\text{P}_2$ levels, thereby regulating channel activity by a local feedback regulation. Whether this local control is a general principle governing the function of other Trp channels, including the 'classical' TrpC channels discussed above, remains to be determined.

A further emerging theme in Ca^{2+} (and other) channel control is that channels located in intracellular membrane compartments are inserted into the plasma membrane by a regulated process quite reminiscent of the GLUT4 glucose transporter insertion into the plasma membrane after insulin stimulation. It has been recently shown that growth factors enhance the insertion of TrpC5 channels from a vesicular pool by a PI3K- and Rac1-mediated mechanism that also involves PIP5KI α (Bezzarides *et al.* 2004). Similarly, PI3K-dependent trafficking of voltage-gated Ca^{2+} channels to the plasma membrane has been recently reported (Viard *et al.* 2004). Intriguingly, a previously described curious stimulatory effect of PLC γ 1 expression on the activity of TrpC3 channels (that oddly did not require the PLC activity of the protein) has also




Figure 5 Multiple steps in the exocytic process. Secretory vesicles containing their cargo become docked to the plasma membrane to form the rapidly releasable vesicular pool by maturation. During maturation, which is an ATP-dependent process, these vesicles become 'primed', that is, ready for release once the Ca^{2+} -induced triggering takes place. Synthesis of phosphoinositides (probably $\text{PtdIns}(4,5)\text{P}_2$) is part of the priming process, and the recruitment of the Ca^{2+} -binding regulatory protein CAPS (which also binds $\text{PtdIns}(4,5)\text{P}_2$) is an essential component of priming. Vesicles are stabilized by the t- and v-snare proteins (syntaxin and VAMP respectively), and they respond to Ca^{2+} increases via synaptotagmin (a protein also capable of phosphoinositide binding), which ultimately triggers the conformational change(s) leading to pore opening and cargo release. Phosphoinositides as well as Ca^{2+} increases act at multiple steps, and production of the lipid both at the membrane and on the surface of the vesicles may take place.

been attributed to enhanced surface expression of the channels (Patterson *et al.* 2002). In this newly discovered process, one of the 1/2 PH-domains of $\text{PLC}\gamma 1$ and a half-PH domain located within the intracellular tails of TrpC channels form an 'intermolecular PH domain' providing phosphoinositide interaction and stabilizing the channel at the plasma membrane (van Rossum *et al.* 2005). These new observations demonstrate that, in addition to the acute regulation of channel activities in the plasma membrane by rapid changes in phosphoinositides, these lipids also control the trafficking and distribution of the channels between the various membranes, adding a new level of complexity to the control of Ca^{2+} (and other ion) fluxes of the cell.

Phosphoinositides and exocytosis

While regulated secretion is one of the key features of many endocrine and neuroendocrine cells, it is also found in other cells such as those of the immune system. This process is also called dense-core vesicle (DCV) exocytosis to distinguish it from the small synaptic vesicle (SV) exocytosis found in presynaptic terminals, although the underlying processes have many common features

(Martin 2003, Wenk & De Camilli 2004). DCVs containing cargo to be secreted undergo maturation that increases their competence to dock, and ultimately fuse, with the plasma membrane when a rapid rise in Ca^{2+} concentration triggers the fusion process. Some of the mature granules under the membrane will dock to the plasma membrane, but these predocked vesicles still have to undergo 'priming' to become the 'readily releasable pool' that is first to be fused upon stimulation (Fig. 5). The prefusion process of 'priming' has been shown to be ATP-dependent and to involve the synthesis of $\text{PtdIns}(4,5)\text{P}_2$ (Eberhard *et al.* 1990, Hay & Martin 1993, Hay *et al.* 1995). Purification of the cytosolic components required for ATP-dependent priming has led to the identification of the PI-transfer protein (Hay & Martin 1993) and a PIP 5-kinase (Hay *et al.* 1995), but the PI4K that is needed for $\text{PtdIns}(4,5)\text{P}_2$ generation has never been identified. It is noteworthy that the first successful cloning of the type II PI4K resulted from the purification of a chromaffin granule-associated PI4K (Barylko *et al.* 2001). Whether this enzyme supports the process of exocytosis is yet to be determined.

The PI3K inhibitor, wortmannin (Wm), potently inhibits the degranulation caused by FcR activation in rat basophilic leukaemia (RBL) cells at concentrations that are consistent with inhibition of PI3K enzymes

(Yano *et al.* 1993). However, in other cells, such as pituitary gonadotrophs (Rao *et al.* 1997), pancreatic β cells (Straub & Sharp 1996) and PC12 cells (Oda *et al.* 1997), higher Wm concentrations were found to inhibit secretion, and, where kinetic analysis was performed, Wm selectively eliminated the sustained phase of secretion without affecting its early rapid phase (Rao *et al.* 1997). These effects of Wm (which were often attributed to MLCK inhibition by early studies) probably result from inhibition of type III PI4K enzymes that are also targets of Wm at the same high concentration range. Wm therefore appears to affect the process of replenishment of the readily releasable pool, at least in some cells. Moreover, a phenylarsine-oxide (PAO)-sensitive PI4K was identified as necessary for secretion in chromaffin cells (Wiedemann *et al.* 1996) and in pancreatic β -cells (Olsen *et al.* 2003). While PAO can have several targets other than PI4Ks, among the latter, PI4 KIII α (another Wm-sensitive PI4K) shows the highest sensitivity to PAO (Balla *et al.* 2002). Recent evidence suggests that the small Ca^{2+} -binding protein, NCS-1, activates the Wm-sensitive PI4 KIII β enzyme and regulates priming and formation of the rapidly releasable vesicular pool in pancreatic β cells (Gromada *et al.* 2005). Similarly, reducing the levels of PI4 KIII β (Waselle *et al.* 2005) or PIP5KI γ (Gong *et al.* 2005, Waselle *et al.* 2005) has been shown to inhibit dense-core vesicle exocytosis. Recent studies have also indicated that PI3KC2 α , by producing PtdIns(3)P, is also involved in neurosecretory granule exocytosis, adding 3-phosphorylated inositides to the list of potential regulators of the priming process (Meunier *et al.* 2005).

The regulatory role of phosphoinositides in synaptic vesicle release and recycling has also been well documented. Both synaptotagmin-1, a phosphoinositide 5-phosphatase (McPherson *et al.* 1996), and PIP5KI γ (Wenk *et al.* 2001) have been shown to associate with synaptic vesicles, and knockout studies have confirmed that these enzymes are essential for normal synaptic functions, and have a role at multiple steps in synaptic vesicle exocytosis and recycling (Cremona *et al.* 1999, Di Paolo *et al.* 2004). Of the PI 4-kinases, PI4KII α was shown to associate with synaptic vesicles (Guo *et al.* 2003). On the other hand, the *Drosophila* homolog of NCS-1, originally named frequenin and described as a major determinant of synaptic development and plasticity (Pongs *et al.* 1993), is a major regulator of the PI4KIII β protein (Weisz *et al.* 2000), as well as its yeast homolog, Pik1p (Hendricks *et al.* 1999). This suggests that there is a link between PI4KIII β function and synaptic transmission. It would not be surprising if both type-II and type-III PI4Ks were found to be important enzymes at distinct steps in the complex process of synaptic vesicle exocytosis and recycling.

Regarding the role of phosphoinositides in exocytosis, two additional questions remain to be answered. The first is whether the phosphoinositides are required on the surface

of the secretory vesicles or at the plasma membrane, or perhaps in both locations. Initial reports have indicated that PtdIns(4,5)P $_2$ is needed on the surface of the vesicles where the presence of the enzymes has been demonstrated (Martin *et al.* 1997). However, studies using the PLC δ 1PH-GFP fusion protein to image PtdIns(4,5)P $_2$ distribution failed to detect this lipid on the vesicular surface (Holz *et al.* 2000), indicating instead a significant increase in PtdIns(4,5)P $_2$ at the contact sites in the plasma membrane associated with the vesicle fusion process (Holz *et al.* 2000, Aoyagi *et al.* 2005). A recent report has established that plasma membrane PtdIns(4,5)P $_2$ levels are a determinant of the size of the readily releasable pool in chromaffin cells (Milosevic *et al.* 2005). Nevertheless, these observations do not rule out additional function(s) of PtdIns(4,5)P $_2$ at the surface of secretory vesicles. Another question relates to the identity of the molecules that are the targets of PtdIns(4,5)P $_2$. CAPS-1 (Ca^{2+} -sensitive activator protein of secretion) has been identified as a crucial factor in DCV exocytosis that acts between docking and fusion (Walent *et al.* 1992). This protein was shown to be a PtdIns(4,5)P $_2$ -binding protein (Loyet *et al.* 1998) whose recruitment to the plasma membrane depended on PtdIns(4,5)P $_2$ levels (Grishanin *et al.* 2004). Interestingly, while CAPS-1 has a central PH domain and PH domains generally serve as lipid-binding modules (Lemmon & Ferguson 2000), the CAPS-1 PH domain is not the principal lipid-binding site of the molecule (Grishanin *et al.* 2004). Other proteins have also been suggested as targets of phosphoinositides in regulated secretion. These include the Ca^{2+} -sensitive SNARE-regulator protein, synaptotagmin, which binds inositides with its C2B domain (Schiavo *et al.* 1996, Bai *et al.* 2004); the Mint proteins (Okamoto & Sudhof 1997); and Rabphilin (Chung *et al.* 1998). Synaptotagmins have key roles in the Ca^{2+} -triggered fusion process (Tucker & Chapman 2002), and their phosphoinositide binding has other functions than vesicle priming, again enforcing the idea that phosphoinositides contribute to the regulation of the secretory process in multiple ways.

Cellular lipid homeostasis and phosphoinositides

Although maintenance of cellular lipid homeostasis is a well-recognized function of all eukaryotic cells, traditionally only some of its aspects are subjects of endocrine research. This is in spite of the fact that lipid metabolism at the level of the whole organism is a central topic of endocrinology. The role or roles of phosphoinositides in the regulation of lipid synthesis and transport are not widely recognized, but recent findings suggest that these processes are also under the control of phosphoinositides. Studies on PH domains that recognize specific inositol lipids have revealed that a special subgroup of PH domains can bind PtdIns(4)P with remarkable specificity

(Dowler *et al.* 2000). Proteins whose PH domains belong to this subgroup are the oxysterol-binding protein (OSBP), the ceramide-transfer protein (CERT) (Hanada *et al.* 2003) and the FAPP1 and FAPP2 proteins (Godi *et al.* 2004). With the exception of the FAPP1 protein, which appears to be a truncated adapter protein, all of these are lipid-binding/transfer proteins. The functions of OSBPs and their related proteins (ORPs) are mostly unknown, but they contain a lipid-binding motif that binds oxysterols or other phospholipids and a PH domain for interaction with membrane phosphoinositides (Lehto & Olkkonen 2003, Olkkonen & Lehto 2004). Oxysterols are among the most potent regulators of the transcription of genes that contribute to cholesterol synthesis and transport (Brown & Goldstein 1974, Olkkonen & Lehto 2004). OSBP overexpression increases cholesterol biosynthesis in chinese hamster ovary (CHO) cells (Lagace *et al.* 1997), and depletion of cholesterol or addition of 25-OH cholesterol promotes the association of OSBP with the Golgi complex (Ridgway *et al.* 1992, Storey *et al.* 1998). Since cholesterol biosynthesis is largely controlled by the SREBP transcription factor, whose nuclear translocation depends on its proteolytic cleavage in the Golgi complex (Brown & Goldstein 1999), the regulation of SREBP vesicular transport from the ER to the Golgi complex is a crucial step in cholesterol homeostasis and one that is likely to be controlled by PI 4-kinase(s). However, little is known about the connection between OSBP proteins and cholesterol homeostasis, and the involvement of phosphoinositides in SREBP trafficking.

More is known about the recently identified CERT protein, which is essential for the transport of ceramide from the site of its synthesis in the ER to the Golgi, where its conversion to sphingomyelin takes place (Perry & Ridgway 2005). CERT also contains a PH domain that binds PtdIns(4)P, and a single-point mutation within the PH domain that eliminates PtdIns(4)P binding is sufficient to render CERT completely dysfunctional (Hanada *et al.* 2003). PtdIns(4)P, therefore, is emerging as an important lipid regulator of the synthesis of sphingomyelin via its participation in the control of ceramide transport between the ER and the Golgi. It is a fascinating question why sufficient amounts of ceramide cannot reach the Golgi via the highly dynamic vesicular transport process that exists between the two organelles. This also suggests that ceramide is excluded from the budding CopII vesicles destined for the Golgi, probably because it is bound to a yet unidentified ER protein. The FAPP2 protein also contains a putative glycolipid-binding domain, but its natural ligand-binding partner is not known. The role (if any) that FAPP2 plays in cellular lipid metabolism, and whether the effects of FAPP2 knockdown on trafficking are related to its lipid-transport function, are also not known (Godi *et al.* 2004, Vieira *et al.* 2005).

Another indication of the importance of PtdIns(4)P in phospholipid synthesis comes from yeast studies.

Synthesis of the aminophospholipid, phosphatidylethanolamine (PE), via decarboxylation of ER-derived phosphatidylserine (PS), takes place either in the mitochondria or in Golgi membranes (Voelker 2005). PS, therefore, has to be transferred to those membranes in order to be decarboxylated, and genetic studies have shown that Stt4p (the yeast homolog of PI4KIII α) is a regulatory component of this process at the ER/Golgi (but not at the mitochondrial) site (Trotter *et al.* 1998). It is not yet known why the Stt4p kinase is needed for lipid transfer or whether it acts at the donor or acceptor membrane site. It also remains to be seen whether a similar regulation of aminophospholipid synthesis by PI 4-kinases or by other phosphoinositides is present in higher organisms.

Sec14p, the yeast PI/PC-TP, is as an essential component of the secretion process from the Golgi in *Saccharomyces cerevisiae*, and also functions in phosphatidylcholine (PC) metabolism and maintains DAG levels in the Golgi (Rout & Bankaitis 2004). There is an intimate relationship between Sec14p and PtdIns(4)P levels within the Golgi, since inactivation of the Sac1p inositol lipid phosphatase (Guo *et al.* 1999) can 'bypass' Sec14p defects (Whitters *et al.* 1993). Mammalian PITP proteins come in various forms: in addition to PITP α and - β , a highly homologous PITP module is found in some of the larger RdgB proteins that are homologs of the *Drosophila* retinal degeneration protein (Vihtelic *et al.* 1993). Two of the PITPs, the small PITP β and large RdgB α 1, are Golgi-localized proteins, and the latter was recently shown to be critical for Golgi morphology and function by controlling DAG levels (Litvak *et al.* 2005). In this regard, the lack of RdgB α 1 in mammalian cells causes a defect similar to that seen in the yeast *SEC14*. The small PITP β appears to be an essential gene in the mouse, but PITP α knockout mice are viable with prominent defects in the transport of re-esterified triglycerides from the ER in enterocytes and a similarly defective lipid handling of hepatocytes (Bankaitis *et al.* 2004). Reduced levels of PITP α are responsible for the early-onset neurodegeneration described in the vibrator mouse (Hamilton *et al.* 1997). PITPs are clearly an exciting group of proteins on the border of phospholipid metabolism and vesicular trafficking, and will surely surprise us with novel functions in the near future.

Nuclear receptors, nuclear signaling and inositol phospholipids

Steroid and thyroid hormones that bind to intracellular receptors and regulate the transcription of their target genes in the nucleus seem to defy the developments of phosphoinositide research. Although a fraction of almost all of the inositide kinases and PLCs have been shown to be present in the nucleus under certain conditions

(Payraastre *et al.* 1992, Manzoli *et al.* 2005), and a separate nuclear phosphoinositide system has long been postulated (Irvine 2003), the link between the latter and the transcriptional apparatus has not been forthcoming. This gap began to narrow when genetic studies in yeast revealed that two inositol polyphosphate (IP) kinases, Ipk1 and Ipk2, are involved in mRNA export, chromatin remodeling and DNA metabolism (Odom *et al.* 2000, Shears 2004). These enzymes are critically important for the synthesis of highly phosphorylated inositols, such as InsP₅ and InsP₆, and also for the production of the pyrophosphorylated PP-InsP₄ and PP-IP₅, but it is not known at this point which (if any) of the highly phosphorylated inositols serves as an active regulatory species. Recent studies showed that the inositol pyrophosphates also regulate telomere length (Saiardi *et al.* 2005, York *et al.* 2005). Most recently, inositol multikinase, the mammalian homolog of Ipk2, was found to function as a wortmannin-insensitive PI 3-kinase in the nucleus. This observation raises the question of whether the lipids or the soluble inositol phosphates are the most important regulatory products of the enzyme (Carroll *et al.* 2004). In a separate line of research, structural characterization of the transcription factor, TFIIF, unveiled the presence of a PH domain in the molecule (Gervais *et al.* 2004). Moreover, an inositide-binding PHD domain was described in the chromatin-associated protein, ING2 (Gozani *et al.* 2003). The transcription factor Tubby, which, when mutated in mice, causes obesity, insulin resistance and sensory deficits (Carroll *et al.* 2004), was also shown to interact within its carboxy-terminal domain with plasma membrane PtdIns(4,5)P₂. Only upon hydrolysis of the lipid does Tubby become free to translocate to the nucleus (Santagata *et al.* 2001). The recent discovery of nuclear orphan receptors that bind phospholipids (Li *et al.* 2005), including PtdIns(3,4,5)P₃, as ligands (Krylova *et al.* 2005) is the latest addition to a list of nuclear proteins potentially regulated by phosphoinositides. It is expected that this area of research will greatly expand in the near future, putting nuclear events on a par with membrane-associated signaling processes as regulatory targets of phosphoinositides.

Concluding remarks

These examples of inositol lipid-regulated processes contributing to the complex cellular responses to hormones and neurotransmitters represent only a limited selection from the multitude of actions of phosphoinositides. The purpose of discussing them in this context was to identify areas of interest and to facilitate the incorporation of phosphoinositides as important topics in endocrine research. Over the last decade, it has become increasingly clear that the molecular elements of the classical messenger cascades, such as receptors, G proteins, ion channels and 'effectors', such as PLCs and A- or G- cyclases, are actively organized

within the cell in the relevant membrane compartments. Phosphoinositides are at least as important in the trafficking and organization of signaling complexes as they are in the rapid generation of intracellular messengers. It is not too difficult to foresee a great expansion of knowledge in this research area and that cellular endocrinology will be an active contributor to, as well as a beneficiary of, this expansion.

Acknowledgements

I thank Dr Kevin J Catt (NICHD, NIH) for his critical reading of the manuscript and helpful suggestions. This research is supported by the Intramural Research Program of the National Institute of Child Health and Human Development of the National Institutes of Health. The author declares that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Abel K, Anderson RA & Shears SB 2001 Phosphatidylinositol and inositol phosphate metabolism. *Journal of Cell Science* **114** 2207–2208.
- Alessi DR, James SR, Downes CP, Holmes AB, Gaffney PR, Reese CB & Cohen P 1997 Characterization of a 3-phosphoinositide-dependent protein kinase which phosphorylates and activates protein kinase B α . *Current Biology* **7** 261–269.
- Aoyagi K, Sugaya T, Umeda M, Yamamoto S, Terakawa S & Takahashi M 2005 The activation of exocytic sites by the formation of phosphatidylinositol 4,5-bisphosphate microdomains at syntaxin clusters. *Journal of Biological Chemistry* **280** 17346–17352.
- Bai J, Tucker WC & Chapman ER 2004 PIP₂ increases the speed of response of synaptotagmin and steers its membrane-penetration activity toward the plasma membrane. *Nature Structural and Molecular Biology* **11** 36–44.
- Balla T 1998 Phosphatidylinositol 4-kinases. *Biochimica Biophysica Acta* **1436** 69–85.
- Balla A, Tuymetova G, Barshishat M, Geiszt M & Balla T 2002 Characterization of type II phosphatidylinositol 4-kinase isoforms reveals association of the enzymes with endosomal vesicular compartments. *Journal of Biological Chemistry* **277** 20041–20050.
- Balla A, Tuymetova G, Tsiomenko A, Varnai P & Balla T 2005 A plasma membrane pool of phosphatidylinositol 4-phosphate is generated by phosphatidylinositol 4-kinase type-III α : studies with the PH domains of the oxysterol binding protein and FAPP1. *Molecular Biology of the Cell* **16** 1282–1295.
- Bankaitis VA, Cortese J, Phillips SE & Alb JG Jr 2004 Phosphatidylinositol transfer protein function in the mouse. *Advances in Enzyme Regulation* **44** 201–218.
- Barylko B, Gerber SH, Binns DD, Grichine N, Khvotchev M, Sudhof TC & Albanesi JP 2001 A novel family of phosphatidylinositol 4-kinases conserved from yeast to humans. *Journal of Biological Chemistry* **276** 7705–7708.
- Berridge MJ 1984 Inositol trisphosphate and diacylglycerol as intracellular messengers. *Biochemical Journal* **220** 345–360.
- Berridge MJ, Downes CP & Hanley MR 1989 Neural and developmental actions of lithium: a unifying hypothesis. *Cell* **59** 411–419.
- Bezprozvanny I, Watras J & Ehrlich BE 1991 Bell-shaped calcium-response curves of Ins(1,4,5)P₃- and calcium-gated

- channels from endoplasmic reticulum of cerebellum. *Nature* **351** 751–754.
- Bezerides VJ, Ramsey IS, Kotecha S, Greka A & Clapham DE 2004 Rapid vesicular translocation and insertion of TRP channels. *Nature Cell Biology* **6** 709–720.
- Bi L, Okabe I, Bernard DJ, Wynshaw-Boris A & Nussbaum RL 1999 Proliferative defect and embryonic lethality in mice homozygous for a deletion in the p110 alpha subunit of phosphoinositide 3-kinase. *Journal of Biological Chemistry* **274** 10963–10968.
- Bi L, Okabe I, Bernard DJ & Nussbaum RL 2002 Early embryonic lethality in mice deficient in the p110 beta catalytic subunit of PI 3-kinase. *Mammalian Genome* **13** 169–172.
- Boehning D & Joseph SK 2000 Direct association of ligand-binding and pore domains in homo- and heterotetrameric inositol 1,4,5-trisphosphate receptors. *EMBO Journal* **19** 5450–5459.
- Bonifacino JS & Weissman AM 1998 Ubiquitin and the control of protein fate in the secretory and endocytic pathways. *Annual Review of Cell and Developmental Biology* **14** 19–57.
- Bonifacino JS & Traub LM 2003 Signals for sorting of transmembrane proteins to endosomes and lysosomes. *Annual Review of Biochemistry* **72** 395–447.
- Bosanac I, Alattia JR, Mal TK, Chan J, Talarico S, Tong FK, Tong KI, Yoshikawa F, Furiuchi T, Iwai M *et al.* 2002 Structure of the inositol 1,4,5-trisphosphate receptor binding core in complex with its ligand. *Nature* **420** 696–700.
- Bosanac I, Yamazaki H, Matsu-ura T, Michikawa T, Mikoshiba K & Ikura M 2005 Crystal structure of the ligand binding suppressor domain of type 1 inositol 1,4,5-trisphosphate receptor. *Molecular Cell* **17** 193–203.
- Brown MS & Goldstein JL 1974 Suppression of 3-hydroxy-3-methylglutaryl coenzyme A reductase activity and inhibition of growth of human fibroblasts by 7-ketocholesterol. *Journal of Biological Chemistry* **249** 7306–7314.
- Brown MS & Goldstein JL 1979 Receptor-mediated endocytosis: insights from the lipoprotein receptor system. *PNAS* **76** 3330–3337.
- Brown MS & Goldstein JL 1999 A proteolytic pathway that controls the cholesterol content of membranes, cells, and blood. *PNAS* **96** 11041–11048.
- Carman CV, Barak LS, Chen C, Liu-Chen L-Y, Onorato JJ, Kennedy SP, Caron MG & Benovic JL 2000 Mutational analysis of Gβγ and phospholipid interaction with G protein-coupled receptor kinase 2. *Journal of Biological Chemistry* **275** 10443–10452.
- Carroll K, Gomez C & Shapiro L 2004 Tubby proteins: the plot thickens. *Nature Reviews. Molecular Cell Biology* **5** 55–63.
- Chemin J, Patel AJ, Duprat F, Lauritzen I, Lazdunski M & Honore E 2005 A phospholipid sensor controls mechanogating of the K⁺ channel TREK-1. *EMBO Journal* **24** 44–53.
- Chong LD, Traynor-Kaplan A, Bokoch GM & Schwartz MA 1994 The small GTP-binding protein Rho regulates a phosphatidylinositol 4-phosphate 5-kinase in mammalian cells. *Cell* **79** 507–513.
- Chuang HH, Prescott ED, Kong H, Shields S, Jordt SE, Basbaum AI, Chao MV & Julius D 2001 Bradykinin and nerve growth factor release the capsaicin receptor from PtdIns(4,5)P₂-mediated inhibition. *Nature* **411** 957–962.
- Chung SH, Song WJ, Kim K, Bednarski JJ, Chen J, Prestwich GD & Holz RW 1998 The C2 domains of Rabphilin3A specifically bind phosphatidylinositol 4,5-bisphosphate containing vesicles in a Ca²⁺-dependent manner. In vitro characteristics and possible significance. *Journal of Biological Chemistry* **273** 10240–10248.
- Cockcroft S 1981 Does phosphatidylinositol breakdown control the Ca²⁺-gating mechanism? *Trends in Pharmacological Sciences* **2** 340–342.
- Cremona O, Di Paolo G, Wenk MR, Luthi A, Kim WT, Takei K, Daniell L, Nemoto Y, Shears SB, Flavell RA *et al.* 1999 Essential role of phosphoinositide metabolism in synaptic vesicle recycling. *Cell* **99** 179–188.
- Csordas G, Thomas AP & Hajnoczky G 1999 Quasi-synaptic calcium signal transmission between endoplasmic reticulum and mitochondria. *EMBO Journal* **18** 96–108.
- da Fonseca PC, Morris SA, Nerou EP, Taylor CW & Morris EP 2003 Domain organization of the type 1 inositol 1,4,5-trisphosphate receptor as revealed by single-particle analysis. *PNAS* **100** 3936–3941.
- Dale LB, Seachrist JL, Babwah AV & Ferguson SS 2004 Regulation of angiotensin II type 1A receptor intracellular retention, degradation, and recycling by Rab5, Rab7, and Rab11 GTPases. *Journal of Biological Chemistry* **279** 13110–13118.
- Daub H, Weiss FU, Wallasch C & Ullrich A 1996 Role of transactivation of the EGF receptor in signalling by G-protein-coupled receptors. *Nature* **379** 557–560.
- Dawson AP & Irvine RF 1984 Inositol(1,4,5)trisphosphate-promoted Ca²⁺ release from microsomal fractions of rat liver. *Biochemical Biophysical Research Communications* **120** 858–864.
- Dikic I 2003 Mechanisms controlling EGF receptor endocytosis and degradation. *Biochemical Society Transactions* **31** 1178–1181.
- Di Paolo G, Moskowitz HS, Gipson K, Wenk MR, Voronov S, Obayashi M, Flavell R, Fitzsimonds RM, Ryan TA & De Camilli P 2004 Impaired PtdIns(4,5)P₂ synthesis in nerve terminals produces defects in synaptic vesicle trafficking. *Nature* **431** 415–422.
- Dowler S, Currie RA, Campbell DG, Deak M, Kular G, Downes CP & Alessi DR 2000 Identification of pleckstrin-homology-domain-containing proteins with novel phosphoinositide-binding specificities. *Biochemical Journal* **351** 19–31.
- Eberhard DA, Cooper CL, Low MG & Holz RW 1990 Evidence that the inositol phospholipids are necessary for exocytosis. *Biochemical Journal* **268** 15–25.
- Franke TF, Kaplan DR, Cantley LC & Toker A 1997 Direct regulation of the Akt protooncogene product by PI(3,4)P₂. *Science* **275** 665–668.
- Fruman DA, Meyers RE & Cantley LC 1998 Phosphoinositide kinases. *Annual Review of Biochemistry* **67** 481–507.
- Fruman DA, Mauvais-Jarvis F, Pollard DA, Yballe CM, Brazil D, Bronson RT, Kahn CR & Cantley LC 2000 Hypoglycaemia, liver necrosis and perinatal death in mice lacking all isoforms of phosphoinositide 3-kinase p85 alpha. *Nature Genetics* **26** 379–382.
- Furuichi T, Yoshikawa S, Miyawaki A, Wada K, Maeda N & Mikoshiba K 1989 Primary structure and functional expression of the inositol 1,4,5-trisphosphate-binding protein P₄₀₀. *Nature* **342** 32–38.
- Gaborik Z & Hunyady L 2004 Intracellular trafficking of hormone receptors. *Trends in Endocrinology and Metabolism* **15** 286–293.
- Gaidarov I, Krupnick JG, Falck JR, Benovic JL & Keen JH 1999 Arrestin function in G protein-coupled receptor endocytosis requires phosphoinositide binding. *EMBO Journal* **18** 871–881.
- Galvan DL, Borrego-Diaz E, Perez PJ & Mignery GA 1999 Subunit oligomerization, and topology of the inositol 1,4,5-trisphosphate receptor. *Journal of Biological Chemistry* **274** 29483–29492.
- Gervais V, Lamour V, Jawhari A, Frindel F, Wasielewski E, Dubaele S, Egly J-M, Thierry J-C, Kieffer B & Poterszman A 2004 TFIIF contains a PH domain involved in DNA nucleotide excision repair. *Nature Structural and Molecular Biology* **11** 616–622.
- Glouchankova L, Krishna UM, Potter BV, Falck JR & Bezprozvanny I 2000 Association of the inositol (1,4,5)-trisphosphate receptor ligand binding site with phosphatidylinositol (4,5)-bisphosphate and adenophostin A. *Molecular Cell Biology Research Communication* **3** 153–158.
- Godi A, Di Campi A, Konstantakopoulos A, Di Tullio G, Alessi DR, Kular GS, Daniele T, Marra P, Lucocq JM & De Matteis MA 2004

- FAPPs control Golgi-to-cell-surface membrane traffic by binding to ARF and PtdIns(4)P. *Nature Cell Biology* **6** 393–404.
- Gong LW, Di Paolo G, Diaz E, Cestra G, Diaz ME, Lindau M, De Camilli P & Toomre D 2005 Phosphatidylinositol phosphate kinase type I gamma regulates dynamics of large dense-core vesicle fusion. *PNAS* **102** 5204–5209.
- Gozani O, Karuman P, Jones DR, Ivanov D, Cha J, Logovskoy AA, Baird CL, Zhu H, Field SJ, Lessnick SL *et al.* 2003 The PHD finger of the chromatin-associated protein ING2 functions as a nuclear phosphoinositide receptor. *Cell* **114** 99–111.
- Grishanin RN, Kowalchuk JA, Klenchin VA, Ann K, Earles CA, Chapman ER, Gerona RR & Martin TF 2004 CAPS acts at a prefusion step in dense-core vesicle exocytosis as a PIP₂ binding protein. *Neuron* **43** 551–562.
- Gromada J, Bark C, Smidt K, Efanov AM, Janson J, Mandic SA, Webb DL, Zhang W, Meister B, Jeromin A *et al.* 2005 Neuronal calcium sensor-1 potentiates glucose-dependent exocytosis in pancreatic beta cells through activation of phosphatidylinositol 4-kinase beta. *PNAS* **102** 10303–10308.
- Guillemette G, Balla T, Baukal AJ & Catt KJ 1988 Characterization of inositol 1,4,5-trisphosphate receptors and calcium mobilization in a hepatic plasma membrane fraction. *Journal of Biological Chemistry* **263** 4541–4548.
- Guo J, Wenk MR, Pellegrini L, Onofri F, Benfenati F & De Camilli P 2003 Phosphatidylinositol 4-kinase type IIalpha is responsible for the phosphatidylinositol 4-kinase activity associated with synaptic vesicles. *PNAS* **100** 3995–4000.
- Guo S, Stolz LE, Lemrow SM & York JD 1999 SAC1-like domains of yeast SAC1, INP52, and INP53 and of human synaptotagmin encode polyphosphoinositide phosphatases. *Journal of Biological Chemistry* **274** 12990–12995.
- Haglund K, Di Fiore PP & Dikic I 2003 Distinct monoubiquitin signals in receptor endocytosis. *Trends in Biochemical Sciences* **28** 598–603.
- Hajnoczky G & Thomas AP 1994 The inositol trisphosphate calcium channel is inactivated by inositol trisphosphate. *Nature* **370** 474–477.
- Hajnoczky G, Robb-Gaspers LD, Seitz MB & Thomas AP 1995 Decoding of cytosolic calcium oscillations in the mitochondria. *Cell* **82** 415–424.
- Hamilton BA, Smith DJ, Mueller KL, Kerrebrock AW, Bronson RT, van Berkel V, Daly MJ, Kruglyak L, Reeve MP, Nemhauser JL *et al.* 1997 The vibrator mutation causes neurodegeneration via reduced expression of P1TP alpha: positional complementation cloning and extragenic suppression. *Neuron* **18** 711–722.
- Hanada K, Kumagai K, Yasuda S, Miura Y, Kawano M, Fukasawa M & Nishijima M 2003 Molecular machinery for non-vesicular trafficking of ceramide. *Nature* **426** 803–809.
- Hardie RC 2003 Regulation of TRP channels via lipid second messengers. *Annual Review of Physiology* **65** 735–759.
- Hay JC & Martin TFJ 1993 Phosphatidylinositol transfer protein required for ATP-dependent priming of Ca²⁺-activated secretion. *Nature* **366** 572–575.
- Hay JC, Fiset PL, Jenkins GH, Fukami K, Takenawa T, Anderson RA & Martin TFJ 1995 ATP-dependent inositol phosphorylation required for Ca²⁺-activated secretion. *Nature* **374** 173–177.
- Hendricks KB, Wang BQ, Schnieders EA & Thorner J 1999 Yeast homologue of neuronal frequenin is a regulator of phosphatidylinositol 4-OH-kinase. *Nature Cell Biology* **1** 234–241.
- Hicke L 2001 A new ticket for entry into budding vesicles – ubiquitin. *Cell* **106** 527–530.
- Hilgemann DW & Ball R 1996 Regulation of cardiac Na⁺, Ca²⁺ exchange and KATP potassium channels by PIP₂. *Science* **273** 956–959.
- Hinchliffe AK, Ciruela A & Irvine RF 1998 PIPKins, their substrates and their products: new functions for old enzymes. *Biochimica Biophysica Acta* **1436** 87–104.
- Holz RW, Hlubek MD, Sorensen SD, Fisher SK, Balla T, Ozaki S, Prestwich GD, Stuenkel EL & Bittner MA 2000 A pleckstrin homology domain specific for PtdIns-4,5-P₂ and fused to green fluorescent protein identifies plasma membrane PtdIns-4,5-P₂ as being important in exocytosis. *Journal of Biological Chemistry* **275** 17878–17885.
- Honda A, Nogami M, Yokozeki T, Yamazaki M, Nakamura H, Watanabe H, Kawamoto K, Nakayama K, Morris AJ, Frohman MA *et al.* 1999 Phosphatidylinositol 4-phosphate 5-kinase alpha is a downstream effector of the small G protein ARF6 in membrane ruffle formation. *Cell* **99** 521–532.
- Horowitz LF, Hirdes W, Suh BC, Hilgemann DW, Mackie K & Hille B 2005 Phospholipase C in living cells: activation, inhibition, Ca²⁺ requirement, and regulation of M current. *Journal of General Physiology* **126** 243–262.
- Hoth M & Penner R 1992 Depletion of intracellular calcium stores activates a calcium current in mast cells. *Nature* **355** 353–356.
- Huang CL, Feng S & Hilgemann DW 1998 Direct activation of inward rectifier potassium channels by PIP₂ and its stabilization by Gbetagamma. *Nature* **391** 803–806.
- Hunyady L, Baukal AJ, Gaborik Z, Olivares-Reyes JA, Bor M, Szaszak M, Lodge R, Catt KJ & Balla T 2002 Differential PI 3-kinase dependence of early and late phases of recycling of the internalized AT1 angiotensin receptor. *Journal of Cell Biology* **157** 1211–1222.
- Irvine RF 2003 Nuclear lipid signalling. *Nature Reviews. Molecular Cell Biology* **4** 349–360.
- Ivetac I, Munday AD, Kisseleva MV, Zhang XM, Luff S, Tiganis T, Whisstock JC, Rowe T, Majerus PW & Mitchell CA 2005 The type Ialpha inositol polyphosphate 4-phosphatase generates and terminates phosphoinositide 3-kinase signals on endosomes and the plasma membrane. *Molecular Biology of the Cell* **16** 2218–2233.
- Jiang Q-X, Thrower EC, Chester DW, Ehrlich BE & Sigworth FJ 2002 Three-dimensional structure of the type 1 inositol 1,4,5-trisphosphate receptor at 2.4 Å resolution. *EMBO Journal* **21** 3575–3581.
- Joly M, Kazlauskas A, Fay FS & Corvera S 1994 Disruption of PDGF receptor trafficking by mutation of its PI-3 kinase binding sites. *Science* **263** 684–687.
- Katzmann DJ, Odorizzi G & Emr SD 2002 Receptor downregulation and multivesicular-body sorting. *Nature Reviews. Molecular Cell Biology* **3** 893–905.
- Kiselyov K, Xu X, Mozhayeva G, Kuo T, Pessah I, Mignieri G, Zhu X, Birnbaumer L & Muallem S 1998 Functional interaction between InsP₃ receptors and store-operated Htrp3 channels. *Nature* **396** 478–482.
- Koreh K & Monaco ME 1986 The relationship of hormone-sensitive and hormone-insensitive phosphatidylinositol to phosphatidylinositol 4,5-bisphosphate in the WRK-1 cell. *Journal of Biological Chemistry* **261** 88–91.
- Krapivinsky G, Gordon EA, Wickman K, Velimirovic B, Krapivinsky L & Clapham DE 1995 The G-protein-gated atrial K⁺ channel IKACH is a heteromultimer of two inwardly rectifying K(+)–channel proteins. *Nature* **374** 135–141.
- Krylova IN, Sablin EP, Moore J, Xu RX, Waitt GM, MacKay JA, Juzumienė D, Bynum JM, Madaus K, Montana V *et al.* 2005 Structural analyses reveal phosphatidylinositols as ligands for the NR5 orphan receptors SF-1 and LRH-1. *Cell* **120** 343–355.
- Lagace TA, Byers DM, Cook HW & Ridgway ND 1997 Altered regulation of cholesterol and cholesteryl ester synthesis in Chinese-hamster ovary cells overexpressing the oxysterol-binding protein is dependent on the pleckstrin homology domain. *Biochemical Journal* **326** 205–213.
- Lee A, Frank DW, Marks MS & Lemmon MA 1999 Dominant-negative inhibition of receptor-mediated endocytosis by a dynamin-1 mutant with a defective pleckstrin homology domain. *Current Biology* **9** 261–264.

- Lee S-J, Xu H, Kang L-W, Amzel LM & Montell C 2003 Light adaptation through phosphoinositide-regulated translocation of *Drosophila* visual arrestin. *Neuron* **39** 121–132.
- Lefkowitz RJ 1993 G protein-coupled receptor kinases. *Cell* **74** 409–412.
- Lehto M & Olkkonen VM 2003 The OSBP-related proteins: a novel protein family involved in vesicle transport, cellular lipid metabolism, and cell signalling. *Biochimica Biophysica Acta* **1631** 1–11.
- Lemmon MA & Ferguson KM 2000 Signal-dependent membrane targeting by pleckstrin homology (PH) domains. *Biochemical Journal* **350** 1–18.
- Leopoldt D, Hanck T, Exner T, Maier U, Wetzker R & Nurnberg B 1998 G $\beta\gamma$ stimulates phosphoinositide 3-kinase- γ by direct interaction with two domains of the catalytic p110 subunit. *Journal of Biological Chemistry* **273** 7024–7029.
- Li Y, Choi M, Cavey G, Daugherty J, Suino K, Kovach A, Bingham NC, Klierer SA & Xu EH 2005 Crystallographic identification and functional characterization of phospholipids as ligands for the orphan nuclear receptor steroidogenic factor-1. *Molecular Cell* **17** 491–502.
- Lintschinger B, Balzer-Geldsetzer M, Baskaran T, Graier WF, Romanin C, Zhu MX & Groschner K 2000 Coassembly of Trp1 and Trp3 proteins generates diacylglycerol- and Ca²⁺-sensitive cation channels. *Journal of Biological Chemistry* **275** 27799–27805.
- Liou J, Kim ML, Heo WD, Jones JT, Myers JW, Ferrell JE Jr & Meyer T 2005 STIM is a Ca²⁺ sensor essential for Ca²⁺-store-depletion-triggered Ca²⁺ influx. *Current Biology* **15** 1235–1241.
- Litvak V, Dahan N, Ramachandran S, Sabanay H & Lev S 2005 Maintenance of the diacylglycerol level in the Golgi apparatus by the Nir2 protein is critical for Golgi secretory function. *Nature Cell Biology* **7** 225–234.
- Liu B, Zhang C & Qin F 2005 Functional recovery from desensitization of vanilloid receptor TRPV1 requires resynthesis of phosphatidylinositol 4,5-bisphosphate. *Journal of Neuroscience* **25** 4835–4843.
- Liu X, Singh BB & Ambudkar IS 2003 TRPC1 is required for functional store-operated Ca²⁺ channels. Role of acidic amino acid residues in the S5–S6 region. *Journal of Biological Chemistry* **278** 11337–11343.
- Lopes CM, Rohacs T, Czizjak G, Balla T, Enyedi P & Logothetis DE 2005 PIP2 hydrolysis underlies agonist-induced inhibition and regulates voltage gating of two-pore domain K⁺ channels. *Journal of Physiology (Lond)* **564** 117–129.
- Loyet KM, Kowalchuk JA, Chaudhary A, Chen J, Prestwich GD & Martin TF 1998 Specific binding of phosphatidylinositol 4,5-bisphosphate to calcium-dependent activator protein for secretion (CAPS), a potential phosphoinositide effector protein for regulated exocytosis. *Journal of Biological Chemistry* **273** 8337–8343.
- Luo J, Field SJ, Lee JY, Engelman JA & Cantley LC 2005 The p85 regulatory subunit of phosphoinositide 3-kinase down-regulates IRS-1 signaling via the formation of a sequestration complex. *Journal of Cell Biology* **170** 455–464.
- Luttrell LM & Lefkowitz RJ 2002 The role of beta-arrestins in the termination and transduction of G-protein-coupled receptor signals. *Journal of Cell Science* **115** 455–465.
- Maeda N, Kawasaki T, Nakade S, Yokota N, Taguchi T, Kasai M & Mikoshiba K 1991 Structural and functional characterization of inositol 1,4,5-trisphosphate receptor channel from mouse cerebellum. *Journal of Biological Chemistry* **266** 1109–1116.
- Majerus PW, Kisseleva MV & Norris FA 1999 The role of phosphatases in inositol signaling reactions. *Journal of Biological Chemistry* **274** 10669–10672.
- Mak DO & Foscett JK 1994 Single-channel inositol 1,4,5-trisphosphate receptor currents revealed by patch clamp of isolated *Xenopus* oocyte nuclei. *Journal of Biological Chemistry* **269** 29375–29378.
- Manzoli L, Martelli AM, Billi AM, Faenza I, Fiume R & Cocco L 2005 Nuclear phospholipase C: involvement in signal transduction. *Progress in Lipid Research* **44** 185–206.
- Marchese A & Benovic JL 2001 Agonist-promoted ubiquitination of the G protein-coupled receptor CXCR4 mediates lysosomal sorting. *Journal of Biological Chemistry* **276** 45509–45512.
- Martin TF 2003 Tuning exocytosis for speed: fast and slow modes. *Biochimica Biophysica Acta* **1641** 157–165.
- Martin TFJ, Loyet KM, Barry VA & Kowalchuk JA 1997 The role of PtdIns(4,5)P₂ in exocytotic membrane fusion. *Biochemical Society Transactions* **25** 1137–1141.
- McPherson PS, Garcia EP, Slepnev VI, David C, Zhang X, Grabs D, Sossin WS, Bauerfeind R, Nemoto Y & De Camilli P 1996 A presynaptic inositol-5-phosphatase. *Nature* **379** 353–357.
- Meunier FA, Osborne SL, Hammond GR, Cooke FT, Parker PJ, Domin J & Schiavo G 2005 Phosphatidylinositol 3-kinase C2 α is essential for ATP-dependent priming of neurosecretory granule exocytosis. *Molecular Biology of the Cell* **16** 4841–4851.
- Michell RH 1982 Is phosphatidylinositol really out of the calcium gate? *Nature* **296** 492–493.
- Mignery GA & Sudhof TC 1990 The ligand binding site and transduction mechanism in the inositol-1,4,5-trisphosphate receptor. *EMBO Journal* **9** 3893–3898.
- Mignery GA, Sudhof TC, Takei K & De Camilli P 1989 Putative receptor for inositol 1,4,5-trisphosphate similar to ryanodine receptor. *Nature* **342** 192–195.
- Mikoshiba K 1993 Inositol 1,4,5-trisphosphate receptor. *Trends in Pharmacological Sciences* **14** 86–89.
- Mikoshiba K 1997 The InsP₃ receptor and intracellular signaling. *Current Opinion in Neurobiology* **7** 339–345.
- Milosevic I, Sorensen JB, Lang T, Krauss M, Nagy G, Haucke V, Jahn R & Neher E 2005 Plasmalemmal phosphatidylinositol-4,5-bisphosphate level regulates the releasable vesicle pool size in chromaffin cells. *Journal of Neuroscience* **25** 2557–2565.
- Mora A, Komander D, Van Aalten DM & Alessi DR 2004 PDK1, the master regulator of AGC kinase signal transduction. *Seminars in Cellular and Developmental Biology* **15** 161–170.
- Mori Y, Wakamori M, Miyakawa T, Hermosura M, Hara Y, Nishida M, Hirose K, Mizushima A, Kurosaki M, Mori E *et al.* 2002 Transient receptor potential 1 regulates capacitative Ca²⁺ entry and Ca²⁺ release from endoplasmic reticulum in B lymphocytes. *Journal of Experimental Medicine* **195** 673–681.
- Naga Prasad SV, Laporte SA, Chamberlain D, Caron MG, Barak L & Rockman HL 2002 Phosphoinositide 3-kinase regulates beta2-adrenergic receptor endocytosis by AP-2 recruitment to the receptor/beta-arrestin complex. *Journal of Cell Biology* **158** 563–575.
- Naga Prasad SV, Jayatilke A, Madamanchi A & Rockman HA 2005 Protein kinase activity of phosphoinositide 3-kinase regulates beta-adrenergic receptor endocytosis. *Nature Cell Biology* **7** 785–796.
- Nakanishi S, Catt KJ & Balla T 1995 A wortmannin-sensitive phosphatidylinositol 4-kinase that regulates hormone-sensitive pools of inositolphospholipids. *PNAS* **92** 5317–5321.
- Nishizuka Y 1988 The molecular heterogeneity of protein kinase C and its implications for cellular regulation. *Nature* **34** 661–665.
- Oda H, Murayama T & Nomura Y 1997 Inhibition of protein kinase C-dependent noradrenaline release by wortmannin in PC12 cells. *Archives in Biochemistry and Biophysics* **337** 96–102.
- Odum AR, Stahlberg A, Wente SR & York JD 2000 A role for nuclear inositol 1,4,5-trisphosphate kinase in transcriptional control. *Science* **287** 2026–2029.
- Odorizzi G, Babst M & Emr SD 1998 Fab1p PtdIns(3)P 5-kinase function essential for protein sorting in the multivesicular body. *Cell* **95** 847–858.
- Okamoto M & Sudhof TC 1997 Mints, Munc18-interacting proteins in synaptic vesicle exocytosis. *Journal of Biological Chemistry* **272** 31459–31464.

- Olkkonen VM & Lehto M 2004 Oxysterols and oxysterol binding proteins: role in lipid metabolism and atherosclerosis. *Annals of Medicine* **36** 562–572.
- Olsen HL, Hoy M, Zhang W, Bertorello AM, Bokvist K, Capito K, Efanov AM, Meister B, Thams P, Yang SN *et al.* 2003 Phosphatidylinositol 4-kinase serves as a metabolic sensor and regulates priming of secretory granules in pancreatic beta cells. *PNAS* **100** 5187–5192.
- Otsu M, Hiles I, Gout I, Fry MJ, Ruiz-Larrea F, Panayotou G, Thompson A, Dhand R, Hsuan JJ *et al.* 1991 Characterization of two 85 kDa proteins that associate with receptor tyrosine kinases, middle-T/pp60^{c-src} complexes, and PI3-kinase. *Cell* **65** 91–104.
- Owen DJ, Collins BM & Evans PR 2004 Adaptors for clathrin coats: structure and function. *Annual Review of Cell and Developmental Biology* **20** 153–191.
- Pacher P & Hajnoczky G 2001 Propagation of the apoptotic signal by mitochondrial waves. *EMBO Journal* **20** 4107–4121.
- Padron D, Wang YJ, Yamamoto M, Yin H & Roth MG 2003 Phosphatidylinositol phosphate 5-kinase Ibeta recruits AP-2 to the plasma membrane and regulates rates of constitutive endocytosis. *Journal of Cell Biology* **162** 693–701.
- Parekh AB & Putney JW Jr 2005 Store-operated calcium channels. *Physiological Reviews* **85** 757–810.
- Parrish WR, Stefan CJ & Emr SD 2004 Essential role for the myotubularin-related phosphatase Ymr1p and the synaptojanin-like phosphatases Sjl2p and Sjl3p in regulation of phosphatidylinositol 3-phosphate in yeast. *Molecular Biology of the Cell* **15** 3567–3579.
- Patel S, Joseph SK & Thomas AP 1999 Molecular properties of inositol 1,4,5-trisphosphate receptors. *Cell Calcium* **25** 247–264.
- Patterson RL, van Rossum DB, Ford DL, Hurt KJ, Bae SS, Suh PG, Kurosaki T, Snyder SH & Gill DL 2002 Phospholipase C-gamma is required for agonist-induced Ca²⁺ entry. *Cell* **111** 529–541.
- Payrastré B, Nievers M, Boonstra J, Breton M, Verkleij AJ & VanBergen en Henegouwen PMP 1992 A differential location of phosphoinositide kinases, diacylglycerol kinase, and phospholipase C in the nuclear matrix. *Journal of Biological Chemistry* **267** 5078–5084.
- Pendaries C, Tronchère H, Plantavid M & Payrastré B 2003 Phosphoinositide signaling disorders in human diseases. *FEBS Letters* **546** 25–31.
- Perry RJ & Ridgway ND 2005 Molecular mechanisms and regulation of ceramide transport. *Biochimica Biophysica Acta* **1734** 220–234.
- Pitcher JA, Freedman NJ & Lefkowitz RJ 1998 G protein-coupled receptor kinases. *Annual Review of Biochemistry* **67** 653–692.
- Pongs O, Lindemeier J, Zhu XR, Theil T, Engelkamp D, Krah-Jentgens I, Lambrecht H-G, Koch KW, Schwemer J, Rivoecchi R *et al.* 1993 Frequentin, a novel calcium-binding protein that modulates synaptic efficacy in the *Drosophila* nervous system. *Neuron* **11** 15–28.
- Putney JW Jr 1986 A model for receptor-regulated calcium entry. *Cell Calcium* **7** 1–12.
- Putney JW Jr 2004 The enigmatic TRPCs: multifunctional cation channels. *Trends in Cell Biology* **14** 282–286.
- Rameh LE, Tolias KF, Duckworth BC & Cantley LC 1997 A new pathway for synthesis of phosphatidylinositol-4,5-bisphosphate. *Nature* **390** 192–196.
- Rao K, Paik WY, Zheng L, Jobin RM, Tomic M, Jiang H, Nakanishi S & Stojilkovic SS 1997 Wortmannin-sensitive and -insensitive steps in calcium-controlled exocytosis in pituitary gonadotrophs: evidence that myosin light chain kinase mediates calcium-dependent and wortmannin-sensitive gonadotropin secretion. *Endocrinology* **138** 1440–1449.
- Rebecchi MJ & Pentyala SN 2000 Structure, function, and control of phosphoinositide-specific phospholipase C. *Physiological Reviews* **80** 1291–1335.
- Revankar CM, Cimino DF, Sklar LA, Arterburn JB & Prossnitz ER 2005 A transmembrane intracellular estrogen receptor mediates rapid cell signaling. *Science* **307** 1625–1630.
- Rhee SG 2001 Regulation of phosphoinositide-specific phospholipase C. *Annual Review of Biochemistry* **70** 281–312.
- Ridgway ND, Dawson PA, Ho YK, Brown MS & Goldstein JL 1992 Translocation of oxysterol binding protein to Golgi apparatus triggered by ligand binding. *Journal of Cell Biology* **116** 307–319.
- Rizzuto R, Brini M, Murgia M & Pozzan T 1993 Microdomains with high Ca²⁺ close to IP₃-sensitive channels that are sensed by neighboring mitochondria. *Science* **262** 744–747.
- Rohacs T, Lopes CM, Michailidis I & Logothetis DE 2005 PI(4,5)P₂ regulates the activation and desensitization of TRPM8 channels through the TRP domain. *Nature Neuroscience* **8** 626–634.
- Roos J, DiGregorio PJ, Yeromin AV, Ohlsen K, Lioudyno M, Zhang S, Safrina O, Kozak JA, Wagner SL, Cahalan MD *et al.* 2005 STIM1, an essential and conserved component of store-operated Ca²⁺ channel function. *Journal of Cell Biology* **169** 435–445.
- Routt SM & Bankaitis VA 2004 Biological functions of phosphatidylinositol transfer proteins. *Biochemistry and Cell Biology* **82** 254–262.
- Runnels LW, Yue LX & Clapham DE 2002 The TRPM7 channel is inactivated by PIP₂ hydrolysis. *Nature Cell Biology* **15** 370–378.
- Saiardi A, Resnick AC, Snowman AM, Wendland B & Snyder SH 2005 Inositol pyrophosphates regulate cell death and telomere length through phosphoinositide 3-kinase-related protein kinases. *PNAS* **102** 1911–1914.
- Saito K, Tolias KF, Saki A, Koon HB, Humphries LA, Scharenberg A, Rawlings DJ, Kinet JP & Carpenter CL 2003 BTK regulates PtdIns-4,5-P₂ synthesis: importance for calcium signaling and PI3K activity. *Immunity* **19** 669–678.
- Santagata S, Boggan TJ, Baird CL, Gomez CA, Zhao J, Shan WS, Myszka DG & Shapiro L 2001 G-protein signaling through tubby proteins. *Science* **292** 2041–2050.
- Sato C, Hamada K, Ogura T, Miyazawa A, Iwasaki K, Hiroaki Y, Tani K, Terauchi A, Fujiyoshi Y & Mikoshiba K 2004 Inositol 1,4,5-trisphosphate receptor contains multiple cavities and L-shaped ligand-binding domains. *Journal of Molecular Biology* **336** 155–164.
- Schiavo G, Gu QM, Prestwich GD, Sollner TH & Rothmann JE 1996 Calcium-dependent switching of the specificity of phosphoinositide binding to synaptotagmin. *PNAS* **93** 13327–13332.
- Seachrist JL, Anborgh PH & Ferguson SS 2000 Beta 2-adrenergic receptor internalization, endosomal sorting, and plasma membrane recycling are regulated by Rab GTPases. *Journal of Biological Chemistry* **275** 27221–27228.
- Shears SB 1998 The versatility of inositol phosphates as cellular signals. *Biochimica Biophysica Acta* **1436** 49–67.
- Shears SB 2004 How versatile are inositol phosphate kinases? *Biochemical Journal* **377** 265–280.
- Shenoy SK, McDonald PH, Kohout TA & Lefkowitz RJ 2001 Regulation of receptor fate by ubiquitination of activated beta 2-adrenergic receptor and beta-arrestin. *Science* **294** 1307–1313.
- Shin HW, Hayashi M, Christoforidis S, Lacas-Gervais S, Hoepfner S, Wenk MR, Modregger J, Uttenweiler-Joseph S, Wilm M, Nystuen A *et al.* 2005 An enzymatic cascade of Rab5 effectors regulates phosphoinositide turnover in the endocytic pathway. *Journal of Cell Biology* **170** 607–618.
- Shisheva A, Sbrissa D & Ikonov O 1999 Cloning, characterization, and expression of a novel Zn²⁺-binding FYVE finger-containing phosphoinositide kinase in insulin-sensitive cells. *Molecular and Cellular Biology* **19** 623–634.
- Shpetner H, Joly M, Hartley D & Corvera S 1996 Potential sites of PI-3 kinase function in the endocytic pathway revealed by the PI-3 kinase inhibitor, wortmannin. *Journal of Cell Biology* **132** 595–605.

- Simoncini T, Hafezi-Moghadam A, Brazil DP, Ley K, Chin WW & Liao JK 2000 Interaction of oestrogen receptor with the regulatory subunit of phosphatidylinositol-3-OH kinase. *Nature* **407** 538–541.
- Sorensen SD, Linsman DA, McEwen EL, Heacock AM & Fisher SK 1998 A role for a wortmannin-sensitive phosphatidylinositol-4-kinase in the endocytosis of muscarinic cholinergic receptors. *Molecular Pharmacology* **53** 827–836.
- Stephens LR, Smrcka AV, Cooke FT, Jackson TR, Sternweis PC & Hawkins PT 1994 A novel phosphoinositide 3 kinase activity in myeloid-derived cells is activated by G protein beta-gamma subunits. *Cell* **77** 83–93.
- Stephens LR, Eguinoa A, Erdjument-Bromage H, Lui M, Cooke F, Coadwell J, Smrcka AS, Thelen M, Cadwallader K, Tempst P *et al.* 1997 The G beta gamma sensitivity of a PI3K is dependent upon a tightly associated adaptor, p101. *Cell* **89** 105–114.
- Storey MK, Byers DM, Cook HW & Ridgway ND 1998 Cholesterol regulates oxysterol binding protein (OSBP) phosphorylation and Golgi localization in Chinese hamster ovary cells: correlation with stimulation of sphingomyelin synthesis by 25-hydroxycholesterol. *Biochemical Journal* **336** 247–256.
- Straub SG & Sharp GW 1996 A wortmannin-sensitive signal transduction pathway is involved in the stimulation of insulin release by vasoactive intestinal polypeptide and pituitary adenylate cyclase-activating polypeptide. *Journal of Biological Chemistry* **271** 1660–1668.
- Strubing C, Krapivinsky G, Krapivinsky L & Clapham DE 2001 TRPC1 and TRPC5 form a novel cation channel in mammalian brain. *Neuron* **29** 645–655.
- Suh BC & Hille B 2002 Recovery from muscarinic modulation of M current channels requires phosphatidylinositol 4,5-bisphosphate synthesis. *Neuron* **35** 507–520.
- Suire S, Coadwell J, Ferguson GJ, Davidson K, Hawkins P & Stephens L 2005 p84, a new Gβγ-activated regulatory subunit of the type IB phosphoinositide 3-kinase p110γ. *Current Biology* **15** 566–570.
- Taylor CW & Laude AJ 2002 IP₃ receptors and their regulation by calmodulin and cytosolic Ca²⁺. *Cell Calcium* **32** 321–334.
- Taylor GS & Dixon JE 2003 PTEN and myotubularins: families of phosphoinositide phosphatases. *Methods in Enzymology* **366** 43–56.
- Terauchi Y, Tsuji Y, Satoh S, Minoura H, Murakami K, Okuno A, Inukai K, Asano T, Kaburagi Y, Ueki K *et al.* 1999 Increased insulin sensitivity and hypoglycaemia in mice lacking the p85 alpha subunit of phosphoinositide 3-kinase. *Nature Genetics* **21** 230–235.
- Thomas GMH, Cunningham E, Fensome A, Ball A, Totty NF, Truong O, Hsuan JJ & Cockcroft S 1993 An essential role of phosphatidylinositol transfer protein in phospholipase C-mediated inositol lipid signaling. *Cell* **74** 919–928.
- Trotter PJ, Wu W-I, Pedretti J, Yates R & Voelker DR 1998 A genetic screen for aminophospholipid transport mutants identifies the phosphatidylinositol 4-kinase, Stt4p, as an essential component in phosphatidylserine metabolism. *Journal of Biological Chemistry* **273** 13189–13196.
- Tucker WC & Chapman ER 2002 Role of synaptotagmin in Ca²⁺-triggered exocytosis. *Biochemical Journal* **366** 1–13.
- van Dam EM, Ten Broeke T, Jansen K, Spijkers P & Stoorvogel W 2002 Endocytosed transferrin receptors recycle via distinct dynamin and phosphatidylinositol 3-kinase-dependent pathways. *Journal of Biological Chemistry* **277** 48876–48883.
- van Rossum DB, Patterson RL, Sharma S, Barrow RK, Kornberg M, Gill DL & Snyder SH 2005 Phospholipase Cgamma1 controls surface expression of TRPC3 through an intermolecular PH domain. *Nature* **434** 99–104.
- Verbsky JW, Chang SC, Wilson MP, Mochizuki Y & Majerus PW 2005 The pathway for the production of inositol hexakisphosphate in human cells. *Journal of Biological Chemistry* **280** 1911–1920.
- Viard P, Butcher AJ, Halet G, Davies A, Nurnberg B, Hebllich F & Dolphin AC 2004 PI3K promotes voltage-dependent calcium channel trafficking to the plasma membrane. *Nature Neuroscience* **7** 939–946.
- Vieira OV, Verkade P & Simons K 2005 FAPP2 is involved in the transport of apical cargo in polarized MDCK cells. *Journal of Cell Biology* **170** 521–526.
- Vihetlic TS, Goeb M, Milligan S, O'Tousa JE & Hyde DR 1993 Localization of *Drosophila* retinal degeneration B, a membrane-associated phosphatidylinositol transfer protein. *Journal of Cell Biology* **122** 1013–1022.
- Voelker DR 2005 Bridging gaps in phospholipid transport. *Trends in Biochemical Sciences* **30** 396–404.
- Walent JH, Porter BW & Martin TF 1992 A novel 145 kd brain cytosolic protein reconstitutes Ca(2+)-regulated secretion in permeable neuroendocrine cells. *Cell* **70** 765–775.
- Wang YJ, Li WH, Wang J, Xu K, Dong P, Luo X & Yin HL 2004 Critical role of PIP5K1γ 87 in InsP3-mediated Ca(2+) signaling. *Journal of Cell Biology* **167** 1005–1010.
- Waselle L, Gerona RR, Vitale N, Martih TF, Bader MF & Regazzi R 2005 Role of phosphoinositide signaling in the control of insulin exocytosis. *Molecular Endocrinology* **19** 3097–3106.
- Watt SA, Kular G, Fleming IN, Downes CP & Lucocq JM 2002 Subcellular localization of phosphatidylinositol 4,5-bisphosphate using the pleckstrin homology domain of phospholipase C delta1. *Biochemical Journal* **363** 657–666.
- Wei YJ, Sun HQ, Yamamoto M, Wlodarski P, Kunii K, Martinez M, Barylko B, Albanesi JP & Yin HL 2002 Type II phosphatidylinositol 4-kinase beta is a cytosolic and peripheral membrane protein that is recruited to the plasma membrane and activated by Rac-GTP. *Journal of Biological Chemistry* **277** 46586–46593.
- Weisz OA, Gibson GA, Leung SM, Roder J & Jeromin A 2000 Overexpression of frequenin, a modulator of phosphatidylinositol 4-kinase, inhibits biosynthetic delivery of an apical protein in polarized madin-darby canine kidney cells. *Journal of Biological Chemistry* **275** 24341–24347.
- Wenk MR & De Camilli P 2004 Protein-lipid interactions and phosphoinositide metabolism in membrane traffic: insights from vesicle recycling in nerve terminals. *PNAS* **101** 8262–8269.
- Wenk MR, Pellegrini L, Klenchin VA, Di Paolo G, Chang S, Daniell L, Arioka M, Martin TF & De Camilli P 2001 PIP kinase Igamm is the major PI(4,5)P₂ synthesizing enzyme at the synapse. *Neuron* **31** 79–88.
- Wes PD, Chevesich J, Jeromin A, Rosenberg C, Stetten G & Montell C 1995 TRPC1, a human homolog of a *Drosophila* store-operated channel. *PNAS* **92** 9652–9656.
- Whitters EA, Cleves AE, McGee TP, Skinner HB & Bankaitis VA 1993 SAC1p is an integral membrane protein that influences the cellular requirement for phospholipid transfer protein function and inositol in yeast. *Journal of Cell Biology* **122** 79–94.
- Wiedemann C, Schäfer T & Burger MM 1996 Chromaffin granule-associated phosphatidylinositol 4-kinase activity is required for stimulated secretion. *EMBO Journal* **15** 2094–2101.
- Wong K, Meyers R & Cantley LC 1997 Subcellular localization of phosphatidylinositol 4-kinase isoforms. *Journal of Biological Chemistry* **272** 13236–13241.
- Xu C, Watras J & Loew LM 2003 Kinetic analysis of receptor-activated phosphoinositide turnover. *Journal of Cell Biology* **161** 779–791.
- Yano H, Nakanishi S, Kimura K, Hanai N, Saitoh Y, Fukui Y, Nonomura Y & Matsuda Y 1993 Inhibition of histamine secretion by wortmannin through the blockade of phosphatidylinositol 3-kinase in RBL-2H3 cells. *Journal of Biological Chemistry* **268** 25846–25856.
- York SJ, Armbruster BN, Greenwell P, Petes TD & York JD 2005 Inositol diphosphate signaling regulates telomere length. *Journal of Biological Chemistry* **280** 4264–4269.

- Zhang H, He C, Yan X, Mirshahi T & Logothetis DE 1999 Activation of inwardly rectifying K⁺ channels by distinct PtdIns(4,5)P₂ interactions. *Nature Cell Biology* **1** 183–188.
- Zhang H, Craciun LC, Mirshahi T, Rohacs T, Lopes CM, Jin T & Logothetis DE 2003 PIP(2) activates KCNQ channels, and its hydrolysis underlies receptor-mediated inhibition of M currents. *Neuron* **37** 963–975.
- Zhang J, Barak LS, Anborgh PH, Laporte SA, Caron MG & Ferguson SS 1999 Cellular trafficking of G protein-coupled receptor/ beta-arrestin endocytic complexes. *Journal of Biological Chemistry* **274** 10999–11006.
- Zhu X, Jiang M, Peyton M, Boulay G, Hurst R, Stefani E & Birnbaumer L 1996 Trp, a novel mammalian gene family essential for agonist-activated capacitative Ca²⁺ entry. *Cell* **85** 661–671.

Received in final form 19 December 2005

Accepted 22 December 2005